arXiv:1109.0898v1 [math.ST] 5 Sep 2011

Detection of a sparse submatrix of a high-dimensional noisy matrix

Butucea, C. *and Ingster, Yu. I.†‡

September 6, 2011

Abstract

We observe a $N \times M$ matrix $Y_{ij} = s_{ij} + \xi_{ij}$ with $\xi_{ij} \sim \mathcal{N}(0,1)$ i.i.d. in i, j, and $s_{ij} \in \mathbb{R}$. We test the null hypothesis $s_{ij} = 0$ for all i, j against the alternative that there exists some submatrix of size $n \times m$ with significant elements in the sense that $s_{ij} \geq a > 0$. We propose test procedures and compute asymptotical detection boundary a in order to have maximal test errors tending to 0 as $M \to \infty,\ N \to \infty,\ p = n/N \to 0,\ q = m/M \to 0$. We prove that this boundary is asymptotically sharp minimax under some additional constraints. Relations with other testing problems are discussed. We propose a testing procedure which adapts to (n, m), which is unknown but belongs to some given set, and compute the adaptive sharp rates. The implementation of our test procedure on synthetic data shows excellent behavior for sparse, not necessarily squared matrices. We extend our sharp minimax results in different directions: first, to Gaussian matrices with unknown variance, second, to matrices of random variables having a distribution from an exponential family (non Gaussian) and, third, to a two-sided alternative for matrices with Gaussian elements.

Keywords: detection of sparse signal, minimax testing, minimax adaptive testing, random matrices, sharp detection bounds.

Mathematics Subject Classification (2000): Primary 62G10, 62G20; Secondary 62C20

1 Introduction

In this paper, we observe a high-dimensional random matrix and we want to test the occurrence of a particular submatrix of much smaller size, which has elements with expected

*University Paris-Est Marne-la-Vallée, 5, bd. Descartes, 77454 Marne-la-Vallée Cedex 2, France and CREST, Timbre J340 3, av. Pierre Larousse, 92240 Malakoff Cedex, France

†Research was partially supported by RFBR grant 11-01-00577 and by the grant NSh–4472.2010.1. The author acknowledges support from the CNRS for his visit to the University Paris-Est Marne-la-Vallée.

‡St.Petersburg State Electrotechnical University, 5, Prof. Popov str., 197376 St.Petersburg, Russia

values larger than some threshold. We assume that the entries of the matrix are independent, identically distributed (i.i.d.) random variables but some underlying phenomenon can increase significantly the expected value of the random variables in the submatrix.

We have the observations that form an $N \times M$ matrix $\mathbf{Y} = \{Y_{ij}\}_{i=1,...,N, j=1,...,M}$:

$$Y_{ij} = s_{ij} + \sigma\xi_{ij}, \quad i = 1, \ldots, N, \quad j = 1, \ldots, M, \tag{1.1}$$

where $\sigma > 0$, $\{\xi_{ij}\}$ are i.i.d. random variables and $s_{ij} \in \mathbb{R}$, for all $i \in \{1, ..., N\}$, $j \in \{1, ..., M\}$. In the first part of the paper, the errors ξ_{ij} are assumed to have Gaussian law with known variance σ^2. Without loss of generality we take $\sigma = 1$ in this case. At the end of the paper, we extend our results in different directions, as discussed later on. We test the null hypothesis that all elements of the matrix $\mathbf{Y}$ are i.i.d., standard Gaussian random variables $\mathcal{N}(0, 1)$, that is

$$H_0 : s_{ij} = 0 \quad \forall \quad i = 1, \ldots, N, \quad j = 1, \ldots, M. \tag{1.2}$$

The alternative under consideration will correspond to $n \times m$-submatrices of sizes $n \in \{1, ..., N\}$, $m \in \{1, ..., M\}$ with large enough entries. Let

$$A \subset \{1, \ldots, N\}, \quad \#(A) = n, \quad B \subset \{1, \ldots, M\}, \quad \#(B) = m, \quad C = A \times B, \tag{1.3}$$

and let $\mathcal{C}_{nm}$ be the collection of all subsets C of the form (1.3). The set $\mathcal{C}_{nm}$ corresponds to the collection of all $n \times m$ submatrices in $N \times M$ matrix. For $a > 0$, which may depend on n, m, N and M. We consider the alternative

$$H_1 : \quad \exists\, C \in \mathcal{C}_{nm} \text{ such that } \; s_{ij} = 0, \text{ if } (i, j) \notin C, \text{ and } s_{ij} \geq a, \text{ if } (i, j) \in C \tag{1.4}$$

(in the Remark 2.1 below we discuss that a slightly larger alternative can be considered). The components of the matrix $\mathbf{Y}$ are independent under the alternative as well. Denote by P_S the probability measure that corresponds to observations (1.1) with matrix $S = \{s_{ij}\}$ and by E_S the expected value with respect to the measure P_S.

Let $\mathcal{S}_{nm,a}$ be the collection of all matrices $S = S_C$ that satisfy (1.4).

We discuss here only right-hand side alternatives, but, obviously, left-hand side alternatives can be treated the same way for variables $-Y_{ij}$ instead of Y_{ij}.

We extend our results to three different setups and sketch the proofs of the results. First, we consider errors having Gaussian distribution with unknown variance σ^2. Then, the errors are assumed to have a distribution which belongs to the exponential family (not necessarily Gaussian). Finally, in the initial case of Gaussian errors with known variance, we consider a two-sided alternative of our test problem.

We are interested here in sparse matrices, i.e. the case when n is much smaller than N and m is much smaller than M.

Sparsity assumptions were introduced for vectors. Estimation as well as hypothesis testing for vectors were thoroughly studied in the literature, see for example Bickel, Ritov and Tsybakov [4] and references therein and Donoho and Jin [5].

In the context of matrices, different sparsity assumptions can be imagined. For example, matrix completion for low rank matrices with the nuclear norm penalization has been studied by Koltchinskii, Lounici and Tsybakov [8]. Other results will be discussed later on.

We study the hypothesis testing problem under a minimax setting. A test is any measurable function of the observations, $\psi = \psi(\{Y_{ij}\})$ taking values in $[0,1]$. For such a test $\psi = \psi(\{Y_{ij}\})$ we denote the first-type error, the second-type error under simple alternative and the maximal second-type error over the set $\mathcal{S}_{nm,a}$ by

$$\alpha(\psi) = E_0\psi, \quad \beta(\psi, S) = E_S(1-\psi), \quad \beta_{nm,a}(\psi) = \sup_{S\in\mathcal{S}_{nm,a}} \beta(\psi, S),$$

respectively. Let the global testing error be the sum of the first and second-type errors:

$$\gamma(\psi, S) = \alpha(\psi) + \beta(\psi, S), \quad \gamma_{nm,a}(\psi) = \sup_{S\in\mathcal{S}_{nm,a}} \gamma(\psi, S) = \alpha(\psi) + \beta_{nm,a}(\psi).$$

We define the minimax second-type error at fixed level $\alpha \in (0,1)$ as

$$\beta_{nm,a,\alpha} = \inf_{\psi:\alpha(\psi)\le\alpha} \beta_{nm,a}(\psi).$$

Similarly, let the minimax global testing risk be

$$\gamma_{nm,a} = \inf_{\psi\in[0,1]} \gamma_{nm,a}(\psi).$$

From now on, we assume in the asymptotics that $N \to \infty,\ M \to \infty$ and $n = n_{NM} \to \infty,\ m = m_{NM} \to \infty$. Other assumptions will be given later.

We suppose that $a > 0$ is unknown. The aim of this paper is to give asymptotically sharp boundaries for minimax testing. That means, we are first interested in the conditions on $a = a_{NM}$ which guarantee distinguishability, i.e., the fact that $\gamma_{nm,a} \to 0$ and $\beta_{nm,a,\alpha} \to 0$ for any $\alpha \in (0,1)$. We construct a testing procedure based on a linear statistic combined with a scan statistic. We prove the upper bounds of the minimax testing risk of this procedure. Second, we describe conditions on a for which we have indistinguishability, i.e., the convergence $\gamma_{nm,a} \to 1$ and $\beta_{nm,a,\alpha} \to 1-\alpha$ for any $\alpha \in (0,1)$. These results are called the lower bounds. The two sets of conditions are complementary and match in rate and constant.

Often the sizes n, m of submatrix are unknown, but we know a set $\mathcal{K}_{NM}$ of couples of indices $(n,m) \in \{1,\dots,N\} \times \{1,\dots,M\}$ containing the true one. Then we consider the "adaptive" problem for the combined alternative $\mathcal{S}_{NM,\mathbf{a}} = \bigcup_{(n,m)\in\mathcal{K}_{NM}} \mathcal{S}_{nm,a_{nm}}$, which corresponds to a collection $\mathbf{a} = \{a_{nm},\ (n,m) \in \mathcal{K}_{NM}\}$. The quantities $\beta_{NM,\mathbf{a},\alpha},\ \gamma_{NM,\mathbf{a}}$

are defined in a similar way as above. We define a testing procedure and check that, if a_{nm} satisfies the conditions for distinguishability uniformly over the collection $\mathbf{a}$, the upper bounds still hold. The adaptive lower bounds hold as an easy consequence of the minimax lower bounds.

The problem of choosing a submatrix in a Gaussian random matrix has been previously studied by Sun and Nobel [9]. They were interested in maximal size submatrices of a matrix with increasing size in two setups. First, they consider the case when the average of the entries of the submatrix is larger than a given threshold and, second, when the entries are well-fitted by a two-way ANOVA matrix in the least-squares sense (i.e., the sum of squares of residuals is smaller than some given threshold).

The algorithm of choosing such submatrices was previously introduced in Shabalin *et al.* [10], who were also interested in finding large average submatrices. This problem is strongly motivated by the research of gene expression in microarray data. In these large matrices it is necessary to recover biclusters, that is associations between sets of samples (rows) and sets of variables (columns). These associations together with clinical and biological information are "a first step in identifying disease subtypes and gene regulatory networks". Many other algorithms for biclustering are discussed and compared on real-data bases concerning breast and lung cancer studies.

Similar problems were considered in Addario-Berry *et al.* [1]. They use the same testing procedures for vectors of random variables, where the alternatives may have various combinatorial structures. In particular, they consider the example of detecting a clique of a certain size in a graph and they compute upper and lower bounds for the Bayesian test error. A bipartite graph of size (N, M) is a graph having edges only between the N vertices of one set to the M vertices of a second set. A biclique is a complete bi-partite subgraph of size (n, m), i.e. a subgraph where all n vertices from the first set are connected to the m vertices from the second set. We consider the problem of detecting a biclique. Our results are sharp minimax and adaptive to the size of the unknown biclique.

The plan of the paper is as follows. In Section 2.1 we give the test procedures. We state the conditions on the detection boundary a such that distinguishability is possible. Under mild additional assumptions we give the conditions on a so that the alternative is indistinguishable from the null hypothesis.

In Section 2.2, we consider the adaptive setup where (n, m) is unknown but belongs to some collection of sequences $\mathcal{K}_{NM}$. We compute the adaptive rates of testing of a slightly modified test procedure.

We include in Section 2.3 discussions of previously studied alternatives: subsets without structure and rectangular submatrices. The first case can be assimilated to detection of a sparse signal in vector observations of length $N \times M$, so the set of alternatives and the detection boundary are much larger than in our case. We summarize well-known results by Ingster [6], Ingster and Suslina [7] and Donoho and Jin [5]. The second case is the

detection of rectangles in the large matrix (connected submatrices), which constitutes a set of alternatives smaller than ours. This case is studied in Arias-Castro *et al.* [2] among these for other geometric shapes of clusters. In order to be self-contained we state and prove sharp upper and lower bounds, for the rectangular clusters.

In Section 3, we perform a numerical study of the procedures that attain the sharp upper bounds. In order to compute the scan statistic a heuristic stochastic algorithm from Shabalin *et al.* [10] is used. The empirical detection boundary is very close to the one predicted by our results.

In Section 4 we give extensions of our results to Gaussian variables of unknown variance σ^2, to non Gaussian matrices with distribution in an exponential family and to two-sided tests for Gaussian matrices, respectively.

Section 5 is mainly concerned with the proof of the lower bounds stated in Section 2.1.3. The Appendix contains the proofs of the other results of the paper.

2 Main results

We denote by $n = n_{NM}$, $m = m_{NM}$ and $a = a_{N,M}$. We recall that asymptotics are taken as

$$N \to \infty, \quad M \to \infty, \quad n \to \infty, \quad m \to \infty.$$

Denote $p = n/N,\ q = m/M$. We suppose, moreover, that

$$p \to 0, \quad q \to 0. \tag{2.1}$$

2.1 Known size of the submatrix

In a minimax setup, we suppose that for each N, M we know n and m.

Let us consider two test procedures, one based on a linear statistic ψ_H^{lin} and the other based on a scan statistic ψ_T^{max}. The final test procedure ψ^* will reject as soon as at least one of them rejects the null hypothesis.

2.1.1 Linear statistic

The first test procedure ψ_H^{lin} is based on the linear statistic

$$t_{lin} = \frac{1}{\sqrt{NM}} \sum_{i,j} Y_{ij}, \quad \psi_H^{lin} = \mathbb{1}_{t_{lin} > H}.$$

One easily gets the following non-asymptotic result.

Proposition 2.1 *We have, for any real number H,*

$$\alpha(\psi_H^{lin}) = \Phi(-H), \quad \beta_{nm,a}(\psi_H^{lin}) \leq \Phi(H - a\sqrt{nmpq}),$$

where Φ denotes the cumulative distribution function of a standard Gaussian random variable.

Proof. Observe that the statistic t_{lin} is standard Gaussian under P_0 which yields the relation for $\alpha(\psi_H^{lin})$. Also $t_{lin} \sim \mathcal{N}(h_{S_C}, 1)$ under P_{S_C}-probability, $S_C \in \mathcal{S}_{nm,a}$, where

$$h_{S_C} = \frac{1}{\sqrt{NM}} \sum_{(i,j)\in C} s_{ij} \geq \frac{anm}{\sqrt{NM}} = a\sqrt{mnpq}.$$

This yields the relation $\beta(\psi_H^{lin}, S) \leq \Phi(H - a\sqrt{mnpq}),\ \forall\, S \in \mathcal{S}_{nm,a}$ and the same inequality for $\beta_{nm,a}(\psi_H^{lin})$. Proposition 2.1 follows. □

2.1.2 Scan statistic

The second test ψ_T^{max} is based on the maximal sum over all submatrices. Put

$$Y_C = \frac{1}{\sqrt{nm}} \sum_{(i,j)\in C} Y_{ij}, \tag{2.2}$$

and

$$t_{max} = \max_{C\in\mathcal{C}_{nm}} Y_C, \quad \psi_T^{max} = \mathbb{1}_{t_{max} > T_{nm}}, \tag{2.3}$$

where $T_{nm} = \sqrt{2\log(G_{nm})},\ G_{nm} = \#(\mathcal{C}_{nm}) = \binom{N}{n}\binom{M}{m}$.

The computation of this statistic is discussed in Section 3.

Proposition 2.2 *Assume (2.1). Then* $\alpha(\psi_T^{max}) \to 0$ *and*

$$\beta_{nm,a}(\psi_T^{max}) \leq \Phi(T_{nm} - a\sqrt{nm}).$$

Proof is given in Section 6.1.

2.1.3 Sharp minimax rates

The following theorem gives necessary conditions for the detection boundary a such that distinguishability holds. The test procedure which attains these bounds is

$$\psi^* = \max\{\psi_H^{lin}, \psi_T^{max}\},$$

for properly chosen H and T_{nm}.

Theorem 2.1 Upper bounds. *Assume (2.1) and let a be such that at least one of the following conditions hold*

$$a^2 nmpq \to \infty \tag{2.4}$$

or

$$\liminf \frac{a^2 nm}{2(n\log(p^{-1}) + m\log(q^{-1}))} > 1. \tag{2.5}$$

Then ψ^* *with* $H \to \infty,\ H < ca\sqrt{nmpq},\ c < 1$, *and with* $T_{nm} = \sqrt{2\log G_{nm}},\ G_{nm} = \binom{N}{n}\cdot\binom{M}{m}$ *is such that* $\gamma_{nm,a}(\psi^*) \to 0$.

Proof By Proposition 2.1, if $H \to \infty$ then $\alpha(\psi_H^{lin}) = \Phi(-H) \to 0$ and if $H < ca\sqrt{nmpq}$,

$$\beta_{nm,a}(\psi_H^{lin}) \leq \Phi((c-1)a\sqrt{nmpq}) \to 0,$$

for $c < 1$. Thus $\gamma_{nm,a}(\psi_H^{lin}) \to 0$.

By Proposition 2.2, $\alpha(\psi_T^{max}) \to 0$. It is not hard to check that under (2.1),

$$\log(G_{nm}) \sim (n\log(p^{-1}) + m\log(q^{-1})). \tag{2.6}$$

Thus, under (2.5),

$$a\sqrt{nm} - T_{nm} = T_{nm}\left(\frac{a\sqrt{nm}(1+o(1))}{\sqrt{2(n\log(p^{-1}) + m\log(q^{-1}))}} - 1\right) \to \infty$$

and this implies that $\gamma_{nm,a}(\psi_T^{max}) \to 0$. □

Theorem 2.2 Lower bounds. *Assume (2.1) and*

$$\frac{\log\log(p^{-1})}{\log(q^{-1})} \to 0, \quad \frac{\log\log(q^{-1})}{\log(p^{-1})} \to 0. \tag{2.7}$$

Moreover, assume that

$$n\log(p^{-1}) \asymp m\log(q^{-1}), \tag{2.8}$$

and that the following two conditions are satisfied:

$$a^2 nmpq \to 0 \tag{2.9}$$

and

$$\limsup \frac{a^2 nm}{2(n\log(p^{-1}) + m\log(q^{-1}))} < 1. \tag{2.10}$$

Then the distinguishability is impossible, i.e., $\gamma_{nm,a} \to 1$ and $\beta_{nm,a,\alpha} \to 1-\alpha$ for any $\alpha \in (0,1)$.

Proof is given in Section 5.

These results for the upper and the lower bounds can be interpreted as follows. Under rather mild conditions (2.1), (2.7) and (2.8), the relations

$$a^2 nmpq \asymp 1, \quad a^2 nm \sim 2(n\log(p^{-1}) + m\log(q^{-1})) \tag{2.11}$$

define a sharp detection boundary in the problem with known (n, m). Note that the detection boundary can be written as

$$a^* = \min\left\{\frac{1}{\sqrt{nmpq}}, \quad \sqrt{\frac{2(n\log(p^{-1}) + m\log(q^{-1}))}{nm}}\right\}.$$

2.2 Adaptation to the size of the submatrix

If the size of the submatrix is unknown, we suppose that, for each N and M, there is a set $\mathcal{K}_{NM}$ containing the unknown pair (n, m), such that

$$\sup_{(n,m)\in\mathcal{K}_{NM}} \left(\frac{1}{n}+\frac{1}{m}+\frac{n}{N}+\frac{m}{M}\right) \to 0,$$

as N, $M \to \infty$. This implies that

$$\sup_{(n,m)\in\mathcal{K}_{NM}} \left(\frac{\log(N)}{n\log(p^{-1})}+\frac{\log(M)}{m\log(q^{-1})}\right) \to 0, \text{ as } N, M \to \infty. \tag{2.12}$$

The adaptive test procedure is $\psi^*_{NM} = \max\{\psi^{lin}_H, \psi^{max}_{NM}\}$, where ψ^{lin}_H is the linear statistic defined in Section 2.1 and ψ^{max}_{NM} is a modified version of ψ^{max}_T defined as follows. Set

$$V_{nm} = \sqrt{2\log(NMG_{nm})}, \quad t_{NM,max} = \max_{(n,m)\in\mathcal{K}_{NM}} \max_{C\in\mathcal{C}_{nm}} Y_C/V_{nm}, \quad \psi^{max}_{NM} = \mathbb{1}_{t_{NM,max}>1}.$$

Proposition 2.3 *Assume (2.1). We have $\alpha(\psi^{max}_{NM}) \to 0$ and*

$$\beta_{nm,\mathbf{a}}(\psi^{max}_{NM}) \le \Phi(\max_{(n,m)\in\mathcal{K}_{NM}} (V_{nm} - a_{nm}\sqrt{nm})).$$

Proof is given in Section 6.1.

Theorem 2.3 *Let the set $\mathcal{K}_{NM}$ be such that condition (2.12) holds.*

Upper bounds. *Let $\mathbf{a} = \mathbf{a}_{NM} = \{a_{nm},\ (n,m) \in \mathcal{K}_{NM}\}$ be such that at least one of the following conditions hold*

$$\min_{(n,m)\in\mathcal{K}_{NM}} a^2_{nm} nmpq \to \infty. \tag{2.13}$$

or

$$\liminf \min_{(n,m)\in\mathcal{K}_{NM}} \frac{a^2_{nm} nm}{2(n\log(p^{-1}) + m\log(q^{-1}))} > 1. \tag{2.14}$$

Then, $\gamma_{NM,\mathbf{a}}(\psi^*_{NM}) \to 0$.

Lower bounds. *Suppose that for each N, M there exists $(n^*, m^*) \in \mathcal{K}_{NM}$ such that*

$$\frac{\log\log(N/n^*)}{\log(M/m^*)} \to 0, \quad \frac{\log\log(M/m^*)}{\log(N/n^*)} \to 0$$

and that

$$n^*\log(N/n^*) \asymp m^*\log(M/m^*),$$

as $N \to \infty$ and $M \to \infty$. Let $\mathbf{a} = \mathbf{a}_{NM} = \{a_{nm},\ (n,m) \in \mathcal{K}_{NM}\}$ be such that

$$a^2_{n^*m^*} n^*m^*p^*q^* \to 0$$

and

$$\limsup \frac{a^2_{n^*m^*} n^*m^*}{2(n^*\log(p^{*-1}) + m^*\log(q^{*-1}))} < 1.$$

Then $\gamma_{NM,\mathbf{a}} \to 1$ and $\beta_{NM,\mathbf{a},\alpha} \to 1-\alpha$ for any $\alpha \in (0,1)$.

Proof. Note that the test ψ_H^{lin} does not depend on n, m. Therefore for distinguishability in the adaptive problem it is sufficient to assume (2.13) (which is a uniform version of (2.4)). We have $\gamma_{NM,\mathbf{a}}(\psi_H^{lin}) \to 0$.

By (2.6), we have

$$\begin{aligned} & \min_{(n,m)\in\mathcal{K}_{NM}} (a_{nm}\sqrt{nm}) - V_{nm}) \\ = & \min_{(n,m)\in\mathcal{K}_{NM}} T_{nm} \left(\frac{a_{nm}\sqrt{nm}(1+o(1))}{\sqrt{2(n\log(p^{-1})+m\log(q^{-1}))}} - \sqrt{1+\frac{\log(NM)}{n\log(p^{-1})+m\log(q^{-1})}} \right) \end{aligned}$$

which goes to infinity under (2.12) and (2.14). Thus, by Proposition 2.3 we have $\gamma_{NM,\mathbf{a}}(\psi_{NM}^{max}) \to 0$.

The lower bounds in the adaptive setup are an obvious consequence of Theorem 2.2. □

Remark 2.1 We can state the alternative hypothesis in a more general form:

$$H_1 : \exists C \in \mathcal{C}_{nm} \text{ such that } s_{ij} = 0, \text{ if } (i,j) \notin C, \text{ and } \sum_{(i,j)\in C} s_{ij} \geq anm.$$

Indeed, our probabilities of error depend on the elements of the submatrix C only through the sum of its elements. Therefore, the previous test procedure will attain the same rates and the same lower bound techniques will give the previous results for this more general test problem.

2.3 Related testing problems

Let us consider two related testing problems under the model (1.1) and the null hypothesis (1.2).

2.3.1 Subsets without structure

Let $\mathcal{D}_k$ consists of all subsets $D \subset \{1,\dots,N\} \times \{1,\dots,M\}$ of cardinality $\#(D) = k$ and let $k = nm$. Let us consider the alternative

$$H_1 : \ \exists\ D \in \mathcal{D}_{nm} \text{ such that } s_{ij} = 0 \text{ if } \ (i,j) \notin D, \text{ and } s_{ij} \geq a \text{ if } \ (i,j) \in D \qquad (2.15)$$

(we do not suppose that the set D is of product structure). Clearly we can consider the matrix $\{Y_{ij}\}$ as a vector of dimension $P = NM$, and the problem is well studied as $P \to \infty$, see Ingster [6], Ingster and Suslina [7], Donoho and Jin [5].

The results are as follows. Let $k = P^{1-\beta}$, $\beta \in (0,1)$. First, let $\beta \leq 1/2$ which corresponds to $k^2 = O(P)$, i.e. $(nm)^2 = O(NM)$. Then the detection boundary is determined by the first condition in (2.11). It means that distinguishability is impossible when $a^2 nmpq \to 0$. On the other hand, if $a^2 nmpq \to \infty$, then distinguishability is provided by the tests of the type ψ_H^{lin}.

Let $\beta \in (1/2, 1)$. Then the detection boundary is determined by the relation

$$a^* \sim \varphi(\beta)\sqrt{\log(P)} = \varphi(\beta)\sqrt{\log(NM)},$$

where

$$\varphi(\beta) = \begin{cases} \sqrt{2\beta - 1}, & 1/2 < \beta \le 3/4, \\ \sqrt{2}(1 - \sqrt{1-\beta}), & 3/4 < \beta < 1, \end{cases} \qquad \beta = 1 - \frac{\log(nm)}{\log(NM)}.$$

This means that, if $\limsup a/(\varphi(\beta)\sqrt{\log(NM)}) < 1$, then distinguishability is impossible, and if $\liminf a/(\varphi(\beta)\sqrt{\log(NM)}) > 1$, then distinguishability is provided by the "high criticism" tests $\psi^{HC} = \mathbb{1}_{\{L_{HC} > H\}}$ based on statistics

$$L(t) = \frac{\sum_{i,j}(\mathbb{1}_{\{Y_{ij} > t\}} - \Phi(-t))}{\sqrt{NM\Phi(t)\Phi(-t)}}, \quad L_{HC} = \max_{t_0 < t} L(t), \quad t_0 > 0,$$

with $H = \sqrt{c \log\log(NM)},\ c > 2$.

2.3.2 Rectangular submatrices

Let $\mathcal{E}_{nm}$ consist of all rectangles of size $n \times m$, i.e., of the sets $E_{kl} = \{k+1, \dots k+n\} \times \{l+1, \dots l+m\},\ 0 \le k \le N-n,\ 0 \le l \le M-m$, and the alternative is of the form

$$H_1 :\ \exists\ E \in \mathcal{E}_{nm} \text{ such that } s_{ij} = 0 \text{ if } \ (i,j) \notin E, \text{ and } s_{ij} \ge a \text{ if } \ (i,j) \in E. \tag{2.16}$$

Similar problems were studied recently in Arias-Castro *et al.* [2] for other related geometrically-shaped clusters. Nevertheless, we give here the proof, since the results of Arias-Castro *et al.* [2] cover only the square matrices in this particular setting.

The detection boundary for (2.16) is determined by

$$a^* \sim \sqrt{\frac{2(\log(p^{-1}) + \log(q^{-1}))}{nm}}.$$

Let us consider the test ψ_Z based on the scan statistic over a particular set of possible rectangles, which is a suitable "grid" on $\mathcal{E}_{nm}$ constructed as follows.

Take $\eta_{nm} = \eta > 0$. Put $n_k = (k-1)n\eta,\ k = 1, ..., K,\ m_l = (l-1)m\eta,\ l = 1, ..., L$, where K, L are such that $N - n(1+\eta) \le n_K \le N - n,\ M - m(1+\eta) \le m_L \le M - m$, which yield $K \sim N/(\eta n),\ L \sim M/(\eta m)$. Put

$$Z_{kl} = \frac{1}{\sqrt{nm}} \sum_{(i,j) \in E_{n_k m_l}} Y_{ij}, \quad Z = \max_{1 \le k \le K,\ 1 \le l \le L} Z_{kl}, \quad \psi_Z = \mathbb{1}_{Z > \sqrt{2\log(KL)}}.$$

In this construction, we considered only $K \times L$ rectangles: $E_{n_k m_l}$ for $k = 1, ..., K$ and $l = 1, ..., L$. Thus, we scan over a number of rectangles which is much smaller than the cardinality of $\mathcal{E}_{nm}$ (for technical reasons) and which is also much larger than the set of non overlapping rectangles (this set would not be large enough).

Theorem 2.4 *Assume (2.1). Then*

Upper bounds. *Let*

$$\liminf \frac{a^2 nm}{2(\log(p^{-1}) + \log(q^{-1}))} > 1,$$

and $\eta = \eta_{nm}$ is taken in such way that $\eta \to 0$, $n\eta \to \infty$, $m\eta \to \infty$, $|\log(\eta)| = o(|\log(pq)|)$. Then $\gamma_{nm,a}(\psi_Z) \to 0$ for the test procedure ψ_Z previously described.

Lower bounds. *Let*

$$\limsup \frac{a^2 nm}{2(\log(p^{-1}) + \log(q^{-1}))} < 1.$$

Then $\gamma_{nm,a} \to 1$, $\beta_{nm,a,\alpha} \to 1 - \alpha$ for any $\alpha \in (0,1)$.

Proof is given in Appendix, Section 6.6.

Note that, the separation rates, i.e., the asymptotics of a that provide distinguishability for the alternative (1.4), are intermediate between the fast separation rates for the alternative (2.16) and the slow rates for the alternative without structure (2.15).

Let us consider the particular case of squared matrices ($N = M$) and squared submatrices ($n = m$) such that $n = N^{1-\beta}$ for some $\beta \in (0,1)$. The sharp asymptotic rates of the detection boundaries can be compared in Table 1.

<table>
<tr><th>Rates</th><th>No structure (2.15)</th><th>Submatrix (1.4)</th><th>Rectangles (2.16)</th></tr>
<tr><td>$\beta \in (0, \frac{1}{3}]$</td><td rowspan="2">$N^{-(1-2\beta)}$</td><td>$N^{-(1-2\beta)}$</td><td rowspan="3">$N^{-(1-\beta)}\sqrt{4\beta\log(N)}$</td></tr>
<tr><td>$\beta \in (\frac{1}{3}, \frac{1}{2}]$</td><td rowspan="2">$N^{-(1-\beta)/2}\sqrt{4\beta\log(N)}$</td></tr>
<tr><td>$\beta \in (\frac{1}{2}, 1)$</td><td>$\varphi(\beta)\sqrt{2\log(N)}$</td></tr>
</table>

Table 1: Table of sharp asymptotic rates of the detection boundary a^* for squared matrices and $n = N^{1-\beta}$

3 Simulations

We have implemented the testing procedure $\psi^* = \max\{\psi_H^{lin}, \psi_T^{max}\}$ on synthetic data. While the linear procedure is rather obvious, the computation of the statistic $t_{max} = \max_{C \in \mathcal{C}_{nm}} Y_C$ is done by using the heuristic algorithm introduced and studied empirically by Shabalin *et al.* [10]. This algorithm is also implemented and studied by Sun and Nobel [9] with good empirical results.

Let us briefly recall this algorithm: we choose randomly a set of n rows out of N. Then, we sum in every column the elements of the previously selected rows. We select

now the columns corresponding to the m largest sums obtained in this way. We sum, next, in every row the elements belonging to the selected columns and select the rows corresponding to the n largest sums. We repeat the algorithm until the sum of elements Y_{ij} of the selected submatrix does not increase anymore. As the procedure can stop at a local maximum, we repeat the procedure K times, where K is large (in our simulation $K = 10000$). We take the maximum value of the outputs. This replication is needed to enforce that with high probability the output approaches the global maximum.

We have simulated matrices of size $N \times M$ of i.i.d. standard Gaussian random variables for $N = M = 200$ and $N = M = 500$.

We calibrated the test statistics ψ_H^{lin} and ψ_T^{max} in such a way that the first-type error occurs with probability $\alpha(\psi^*) \leq 1\%$. This calibration is done by using the Gaussian quantile $H = 2.3262$ for ψ_H^{lin} and the empirical quantile (out of 100 samples) for ψ_T^{max}.

Then, we have added the value $a > 0$ to the elements of the upper left submatrix of size $n \times m$. From resulting observations, we compute $\psi^* = \max\{\psi_H^{lin}, \psi_T^{max}\}$. We repeat the test $L = 100$ times and average the values of the test procedure ψ^*. Denote by $\bar{\psi}^*$ this average and note that $1 - \bar{\psi}^*$ estimates the second-type error probability.

We plot the estimated second-type error probabilities for different values of a in the neighborhood of the detection boundary predicted by our theorems, for different values of n and m. The results in Figure 1 correspond to $N = M = 200$, while in Figure 2 to $N = M = 500$.



Figure 1: Estimated second-type error probability for fixed $\alpha = 1\%$, detection boundary a^*, $N = M = 200$

Figures 1 and 2 show that the empirical detection boundary is very close to a^* which is predicted by out theoretical results. Indeed, the second-type error probability is close to 0.5 at some point close to a^*. The plots also show very fast decay of this probability on a small vicinity of a^*. This means that the test is very powerfull above a small neighbourhood of the detection boundary a^*. Note also that, for fixed N and M, a^* decreases to 0 as n and m increase.



Figure 2: Estimated second-type error probability for fixed $\alpha = 1\%$, detection boundary a^*, $N = M = 500$

4 Extensions

We extend our results in different directions. First, we consider matrices of i.i.d. random variables having Gaussian law with unknown variance σ, second, random variables having a distribution belonging to the exponential family (not necessarily Gaussian) and, third, test problem with two-sided alternative for the Gaussian matrices.

4.1 Extension to Gaussian variables with unknown variance

Sharp results in Theorems 2.1 and 2.2 still hold if the random variables Y_{ij} have unknown variance σ, under a mild additional assumption. We sketch here the test procedure and proof of the upper bounds.

We estimate the unknown variance σ^2 of our data by $\hat{\sigma}^2$, where

$$\hat{\sigma}^2 = \frac{1}{NM} \sum_{i,j} Y_{ij}^2.$$

This estimator is unbiased under the null hypothesis, but biased under the alternative.

We replace Y_{ij} by $Y_{ij}/\hat{\sigma}$ in the test procedure ψ^*. We denote by $\hat{t}_{lin} = t_{lin}/\hat{\sigma}$, $\hat{t}_{max} = t_{max}/\hat{\sigma}$ and put

$$\hat{\psi}^* = \max\{\mathbb{1}_{\hat{t}_{lin} > H}, \mathbb{1}_{\hat{t}_{max} > T_{nm}}\}.$$

Theorem 4.1 *Assume (2.1). We suppose that*

$$\max_{i,j} \frac{s_{ij}}{\sigma\sqrt{NM}} = o(1). \tag{4.1}$$

If a is such that one of the following conditions hold

$$\frac{a}{\sigma}\sqrt{nmpq} \to \infty \text{ or } \liminf \frac{a\sqrt{nm}}{\sqrt{2\sigma(n\log(p^{-1}) + m\log(q^{-1}))}} > 1$$

then ψ^*, with $H \to \infty$ such that $H < O((a^2 nmpq/\sigma^2)^{1/4})$ and with $T_{nm} = \sqrt{(2+\delta)\log(\binom{N}{n}\binom{M}{m})}$ for some $\delta > 0$ small enough, is such that $\gamma_{nm,a}(\psi^*) \to 0$.

Proof is given in Section 6.7.

4.2 Extension to general law from an exponential family

In many applications, we do not have Gaussian observations. Instead, we have observations X_{ij}, i.i.d. with probability density $g_{\theta_{ij}}$ from an exponential family, for all $i = 1, ..., N$, and $j = 1, ..., M$. We explain here how to use the previous testing procedures in order to deal with such setups and check that results similar to the case of Gaussian variables hold in this case.

We assume that the laws belong to an exponential family in the general form

$$g_\theta(x) = e^{\eta(\theta)\cdot T(x) - C(\theta)} h(x),\ \theta \in \Theta, \tag{4.2}$$

for the dominating measure μ, where η is supposed 2 times continuously differentiable and strictly increasing on Θ, i.e. $\eta'(\theta) > 0$. Recall that

$$C(\theta) = \log\left(\int e^{\eta(\theta)\cdot T(x)} h(x)\mu(dx)\right).$$

We consider a point θ^0 interior to Θ and test the following hypotheses:

$$H_0 : \theta_{ij} = \theta^0, \text{ for all } i = 1, ..., N,\ j = 1, ..., M$$

against the alternative

$$H_1 : \exists C \in \mathcal{C}_{nm} \text{ such that } \theta_{ij} = \theta^0 \text{ if } (i,j) \notin C \text{ and } \theta_{ij} - \theta^0 \geq d \text{ if } (i,j) \in C. \tag{4.3}$$

Let us write model (4.2) in the canonical form: $g_\eta(x) = \exp(\eta \cdot T(x) - B(\eta))h(x)$, where $\eta = \eta(\theta)$ and $B(\eta) = \log\left(\int e^{\eta T(x)} h(x)\mu(dx)\right)$. Note that, if $\Delta\theta := \theta - \theta^0$ is small, then $\Delta\eta := \eta - \eta^0 \sim \eta'(\theta^0)\Delta\theta$ and $\Delta\eta > 0$.

Now, let us change the variable and put $Y = (T(X) - m^0)/\sigma^0$, with $m^0 = E_{\eta^0}(T(X))$ and $\sigma^0 = \sqrt{Var_{\eta^0}(T(X))}$, having density

$$f_s(y) = \exp(s \cdot y - A(s))h(y),$$

where $s = \eta\sigma_0$ and $A(s) = B(s/\sigma^0) - sm_0/\sigma_0$. Here, we have $A'(s^0) = 0$, $A''(s^0) = 1$ and

$$A(s^0 + \Delta s) - A(s^0) \sim \frac{(\Delta s)^2}{2} \text{ and } A'(s^0 + \Delta s) \sim (\Delta s), \text{ as } \Delta s \to 0. \tag{4.4}$$

The original problem corresponds to testing the null hypothesis $s_{ij} = s^0$ for all (i, j) against the alternative that, for some submatrix $C \in \mathcal{C}_{nm}$, $s_{ij} - s^0 \geq \Delta s = \sigma^0 \Delta\eta$ if and only if $(i, j) \in C$. Set $a = \Delta s$.

We have the following results for exponential models.

Theorem 4.2 *Assume (2.1). We suppose that*

$$\frac{\log(p^{-1})}{m} + \frac{\log(q^{-1})}{n} \to 0. \tag{4.5}$$

Upper bounds. *If a is such that one of the following conditions hold*

$$A'(s^0 + a)\sqrt{nmpq} \to \infty \text{ or } \liminf \frac{A'(s^0 + a)\sqrt{nm}}{\sqrt{2(n\log(p^{-1}) + m\log(q^{-1}))}} > 1$$

then ψ^*, with $H \to \infty$ such that $H < cA'(s^0 + a)\sqrt{nmpq}$ for some $0 < c < 1$ and with $T_{nm} = \sqrt{(2+\delta)\log(\binom{N}{n}\binom{M}{m})}$ for some $\delta > 0$ small enough, is such that $\gamma_{nm,a}(\psi^*) \to 0$.

Lower bounds. *Assume, moreover, that conditions (2.7) and (2.8) hold. If a is such that the following two conditions are satisfied:*

$$a\sqrt{nmpq} \to 0 \text{ and } \limsup \frac{a\sqrt{nm}}{\sqrt{2(n\log(p^{-1}) + m\log(q^{-1}))}} < 1,$$

then $\gamma_{nm,a} \to 1$ and $\beta_{nm,a,\alpha} \to 1 - \alpha$ for any $\alpha \in (0, 1)$.

Proof of the upper bounds is given in Section 6.8.

The proof of the lower bounds uses the relation (4.4) and follows exactly the same lines as the proof of Theorem 2.2 in Section 2.1.3 except that we have to consider $T_{kl}^2 \sim (2+\delta)(k\log(p^{-1}) + l\log(q^{-1}))$ for some small $\delta > 0$ instead of thresholds in (5.3). □

Under the assumption (4.5), the detection boundary $a^* \to 0$. Therefore,

$$A'(s^0 + a^*) \sim a^* \sim (\eta - \eta^0)\sigma^0 \sim \eta'(\theta^0)\sigma^0 d^*,$$

as $d^* \to 0$. It is well known that the Fisher information at θ^0 in model (4.2) is $I(\theta_0) = (\sigma^0\eta'(\theta^0))^2$. In this way, we deduce the sharp asymptotic detection boundary for alternative (4.3) from Theorem 4.2: $d^* = a^*/\sqrt{I(\theta^0)}$. Examples of such calculations for most popular probability distributions in the exponential family are given in Table 2.

Probability law	η	m^0	σ^0	$\sqrt{I(\theta^0)} = \sigma^0\eta'(\theta^0)$
Poisson(θ), $\theta > 0$	$\log(\theta)$	θ^0	$\sqrt{\theta^0}$	$(\theta^0)^{-1/2}$
Ber(θ), $0 < \theta < 1$)	$\log(\frac{\theta}{1-\theta})$	θ^0	$\sqrt{\theta^0(1-\theta^0)}$	$(\theta^0(1-\theta^0))^{-1/2}$
Exp(θ), $\theta > 0$	$-\theta^{-1}$	θ^0	θ^0	$(\theta^0)^{-1}$
$N(0, \theta^2)$, $\theta > 0$	$-\frac{1}{2\theta^2}$	$(\theta^0)^2$	$2(\theta^0)^2$	$2(\theta^0)^{-1}$

Table 2: Examples of calculations for testing in general exponential families

4.3 Extension to two-sided alternative

Let us consider model (1.1) and the same null hypothesis (1.2), against the two-sided alternative:

$$H_1 : \exists C \in \mathcal{C}_{nm} \text{ such that } s_{ij} = 0, \text{ if } (i,j) \notin C \text{ and } |s_{ij}| \geq a, \text{ if } (i,j) \in C.$$

Let us consider the following test procedures

$$z_{lin} = \frac{1}{\sqrt{2NM}} \sum_{i,j} (Y_{ij}^2 - 1) \text{ and } \psi_{lin}^z = \mathbb{1}_{z_{lin} > H}$$

and

$$z_{max} = \max_{C \in \mathcal{C}_{nm}} Z_C, \text{ where } Z_C = \frac{1}{\sqrt{2nm}} \sum_{(i,j) \in C} (Y_{ij}^2 - 1), \text{ and } \psi_{max}^z = \mathbb{1}_{z_{max} > T}.$$

Theorem 4.3 *Assume (2.1). We suppose that (4.5) holds.*

Upper bounds. *If a is such that one of the following conditions hold*

$$a^2\sqrt{nmpq} \to \infty \text{ or } \liminf \frac{a^2\sqrt{nm}}{2\sqrt{n\log(p^{-1}) + m\log(q^{-1})}} > 1$$

then $\psi^z = \max\{\psi_{lin}^z, \psi_{max}^z\}$ with $H \to \infty$ such that $H < ca^2/2\sqrt{nmpq}$ for some $0 < c < 1$ and with $T_{nm} = \sqrt{(2+\delta)\log(\binom{N}{n}\binom{M}{m})}$ for some $\delta > 0$ small enough, is such that $\gamma_{nm,a}(\psi^z) \to 0$.

Lower bounds. *Assume, moreover, that conditions (2.7) and (2.8) hold. If a is such that the following two conditions are satisfied:*

$$a^2\sqrt{nmpq} \to 0 \text{ and } \limsup \frac{a^2\sqrt{nm}}{2\sqrt{n\log(p^{-1}) + m\log(q^{-1})}} < 1,$$

then $\gamma_{nm,a} \to 1$ and $\beta_{nm,a,\alpha} \to 1 - \alpha$ for any $\alpha \in (0,1)$.

Proof is given in Section 6.9.

5 Proof of Theorem 2.2

In the first part, we give the proof of the theorem and the other parts of this section are dedicated to proofs of intermediate results. More lemmas are in the Appendix.

5.1 Prior and truncated likelihood ratio

Let $S_C = \{s_{ij}\}$ be the matrix such that $s_{ij} = 0,\ (i,j) \notin C,\ s_{ij} = a,\ (i,j) \in C$. Let us consider the prior on the set of matrices:

$$\pi = G^{-1} \sum_{C \in \mathcal{C}_{nm}} \delta_{S_C}, \quad G = \#(\mathcal{C}_{nm}),$$

and let P_π be the mixture of likelihoods $P_\pi = G^{-1} \sum_{C \in \mathcal{C}_{nm}} P_{S_C}$. Let us consider the likelihood ratio

$$L_\pi(Y) = \frac{dP_\pi}{dP_0}(Y) = G^{-1} \sum_{C \in \mathcal{C}_{nm}} \frac{dP_{S_C}}{dP_0}(Y) = G^{-1} \sum_{C \in \mathcal{C}_{nm}} \exp(-b^2/2 + bY_C),$$

here and below we set $b^2 \triangleq a^2 nm$, and, for submatrix C of the size $n \times m$, the statistics Y_C are defined by (2.2). Since $\pi(\mathcal{S}_{nm}) = 1$, in order to obtain indistinguishability: $\gamma_{nm,a} \to 1,\ \beta_{nm,a,\alpha} \to 1 - \alpha,\ \forall\ \alpha \in (0,1)$, it suffices to show

$$L_\pi(Y) \to 1 \quad \text{in } P_0 - \text{probability}. \tag{5.1}$$

Indeed,

$$\begin{aligned}
\gamma_{nm,a} &= \inf_{\psi \in [0,1]} \sup_{S \in \mathcal{S}_{nm,a}} (\alpha(\psi) + \beta(\psi, S)) \\
&\geq \inf_{\psi \in [0,1]} \frac{1}{G} \sum_{S \in \mathcal{S}_{nm,a}} \left(E_0(\psi(Y)) + E_0 \left[(1 - \psi(Y)) \frac{dP_{S_C}}{dP_0}(Y) \right] \right) \\
&\geq \inf_{\psi \in [0,1]} \left(E_0(\psi(Y)) + E_0 \left[(1 - \psi(Y)) L_\pi(Y) \right] \right) \\
&\geq E_0(\psi^*(Y)) + E_0 \left[(1 - \psi^*(Y)) L_\pi(Y) \right],
\end{aligned}$$

where $\psi^*(Y) = \mathbb{1}_{L_\pi(Y) > 1}$ is the likelihood ratio test. Therefore (5.1) implies by Fatou lemma that

$$\liminf \gamma_{nm,a} \geq E_0 \left[\liminf \left(E_0(\psi^*(Y)) + E_0[(1 - \psi^*(Y)) L_\pi(Y)] \right) \right],$$

i.e. $\gamma_{nm,a} \to 1$. It is easy to deduce that $\beta_{nm,a,\alpha} \to 1 - \alpha$.

Let us replace the statistics $L_\pi(Y)$ by their truncated version

$$\tilde{L}_\pi(Y) = G^{-1} \sum_{C \in \mathcal{C}_{nm}} \frac{dP_{S_C}}{dP_0}(Y) \mathbb{1}_{Z_C},$$

where the events Z_C are determined as follows. Set

$$T_{kl} = \sqrt{2(\log(G_{kl}) + \log(nm))} \to \infty.$$

Take small $\delta_1 > 0$ (which will be specified later) and set $k_0 = \delta_1 n,\ l_0 = \delta_1 m$. Let $\mathcal{C}_{kl,C} = \{V \in \mathcal{C}_{kl} :\ V \subset C\}$ be the sub-matrices of $C \in \mathcal{C}_{nm}$ which are in $\mathcal{C}_{kl}$. Then we set

$$Z_C = \bigcap_{k_0 \leq k \leq n,\ l_0 \leq l \leq m} \bigcap_{V \in \mathcal{C}_{kl,C}} \{Y_V \leq T_{kl}\}. \tag{5.2}$$

By (2.6), under conditions on k, l in (5.2) (and similarly to the equivalent of T^2_{nm}) we have

$$T^2_{kl} \sim 2(k \log(p^{-1}) + l \log(q^{-1})). \tag{5.3}$$

Indeed, when looking at second-order moments of the likelihood ratio $L_\pi(Y)$ a large contribution comes from overlapping submatrices inducing correlated random variables Y_{C_1} and Y_{C_2}. Then V is the submatrix where C_1 and C_2 overlap. Our idea is to truncate Y_V for those matrices which overlap significantly ($\delta_1 n \le k \le n$ and $\delta_1 m \le l \le m$), hence the likelihood $\tilde{L}_\pi(Y)$.

Proposition 5.1 *Set $Z_{nm} = \bigcap_{C \in \mathcal{C}_{nm}} Z_C$. Then, under the assumptions of Theorem 2.2, $P_0(Z_{nm}) \to 1$.*

Proof is given in Appendix, Section 6.2.

Proposition 5.1 yields

$$P_0\left(L_\pi(Y) = \tilde{L}_\pi(Y)\right) \to 1,$$

and in place of (5.1) it suffices to check that

$$\tilde{L}_\pi(Y) \to 1 \quad \text{in } P_0 - \text{probability}. \tag{5.4}$$

In order to get (5.4) it suffices to verify two relations:

Proposition 5.2 *Under the assumptions of Theorem 2.2, we have*

$$E_0(\tilde{L}_\pi) \to 1.$$

Proof is given in Appendix, Section 6.3.

This Proposition together with the fact that

$$E_0(\tilde{L}^2_\pi) \le 1 + o(1) \tag{5.5}$$

imply that

$$E_0(\tilde{L}_\pi - 1)^2 = \left(E_0(\tilde{L}^2_\pi) - 1\right) - 2\left(E_0(\tilde{L}_\pi) - 1\right) \le o(1)$$

which ends the proof of the theorem.

The remaining part of this section is devoted to obtaining the relation (5.5).

5.2 Second order moment of the truncated likelihood ratio

Let us prove the relation (5.5). We have

$$E_0(\tilde{L}^2_\pi) = G^{-2} \sum_{C_1 \in \mathcal{C}_{nm}, C_2 \in \mathcal{C}_{nm}} E_0\left(\exp(-b^2 + b(Y_{C_1} + Y_{C_2}))\mathbb{1}_{\{Z_{C_1} \cap Z_{C_2}\}}\right). \tag{5.6}$$

Let $C_1 = A_1 \times B_1,\ C_2 = A_2 \times B_2$ and set

$$k = \#(A_1 \cap A_2),\ l = \#(B_1 \cap B_2), \quad V = (A_1 \cap A_2) \times (B_1 \cap B_2), \quad \rho_{kl} = kl/nm.$$

Denote

$$g(k,l) = E_0\left(\exp(-b^2 + b(Y_{C_1} + Y_{C_2}))\mathbb{1}_{\{Z_{C_1}\cap Z_{C_2}\}}\right)$$

In view of symmetry we can fix C_1, say $C_1 = \{1,\dots n\}\times\{1,\dots m\}$, and take the sums over C_2 only. Therefore we have

$$E_0(\tilde{L}_\pi^2) = G^{-1}\sum_{C_2\in\mathcal{C}_{nm}} g(k,l). \tag{5.7}$$

Set $z_{kl}^2 = a^2kl,\ \rho_{kl} = kl/nm$.

Lemma 5.1 *The following inequalities hold true.*

(1) We have

$$g(k,l) \le E_0\left(\exp(-b^2 + b(Y_{C_1} + Y_{C_2}))\right) = \exp(z_{kl}^2) \triangleq g_1(k,l). \tag{5.8}$$

(2) Let $b \ge T_{nm}/(1+\rho_{kl})$. Then

$$\begin{aligned} g(k,l) &\le E_0\left(\exp(-b^2 + b(Y_{C_1} + Y_{C_2}))\,\mathbb{1}_{\{Y_{C_1}\le T_{nm},\, Y_{C_2}\le T_{nm}\}}\right) \\ &\le \exp\left(-(T_{nm}-b)^2 + \frac{\rho_{kl}T_{nm}^2}{1+\rho_{kl}}\right) \triangleq g_2(k,l). \end{aligned} \tag{5.9}$$

(3) Let $k \ge \delta_1 n,\ l \ge \delta_1 m$, and $T_{kl} \le 2z_{kl}$. Then

$$\begin{aligned} g(k,l) &\le E_0\left(\exp(-b^2 + b(Y_{C_1} + Y_{C_2}))\,\mathbb{1}_{\{Y_{u_{kl}}\le T_{kl}\}}\right) \\ &= \exp(T_{kl}^2/2 - (T_{kl} - z_{kl})^2) \triangleq g_3(k,l). \end{aligned} \tag{5.10}$$

Proof of Lemma 5.1 is given in Appendix, Section 6.4.

5.2.1 From hypergeometric to binomial distributions

Observe that the right-hand side of (5.7) is the expectation of $g(X_1, X_2)$ over X_1, X_2 which are independent and having hypergeometric distributions $\mathcal{HG}_1 = \mathcal{HG}(N,n,n),\ \mathcal{HG}_2 = \mathcal{HG}(M,m,m)$ respectively, i.e.,

$$G^{-1}\sum_{C_2\in\mathcal{C}_{nm}} g(k,l) = E_{\mathcal{HG}_1\times\mathcal{HG}_2} g(X_1,X_2). \tag{5.11}$$

Let us compare random variables X having hypergeometric distributions $\mathcal{HG} = \mathcal{HG}(N,n,n)$ and binomial distribution $Bin = Bin(n,\tilde{p}),\ \tilde{p} = n/(N-n)$.

Lemma 5.2 *Under binomial distribution, X is stochastically larger, than under hypergeometric distributions, i.e. for any $x\in\mathbb{R}$,*

$$P_{\mathcal{HG}}(X \ge x) \le P_{Bin}(X \ge x).$$

This yields, for any non-decreasing function $g(x)$,

$$E_{\mathcal{HG}}(g(X)) \le E_{Bin}(g(X)).$$

Proof. The first claim corresponds to Lemma 3 in Arias-Castro *et al.* [3]. The second claim follows from the Abel's transform of the series for the expectation. □

Let $P_{n,p}(k) = P_{Bin}(X = k)$, for some integer k, where X has binomial $Bin(n,p)$ distribution, and similarly $P_{N,n,n}(k) = P_{\mathcal{HG}}(X = k)$ for hypergeometric distributions $\mathcal{HG}(N,n,n)$ of X.

Lemma 5.3 *Let* $n \to \infty,\ p \to 0,\ \ p > 0,\ k \geq n/r(p)$ *where* $r(p) \geq 1$ *for* $p > 0$ *small enough, and* $\log(r(p)) = o(\log(p^{-1}))$. *Then*

$$\log(P_{n,p}(k)) \leq k \log(p)(1 + o(1)).$$

Proof is given in Appendix, Section 6.5

Since $\tilde{p} \sim p$, these imply the following lemma.

Lemma 5.4 *Under assumption of Lemma 5.3,*

$$\log(P_{N,n,n}(k)) \leq k \log(p)(1 + o(1)).$$

Proof. In view of Lemmas 5.2, 5.3 we have

$$P_{N,n,n}(k) \leq P_{\mathcal{HG}}(Z \geq k) \leq P_{Bin}(Z \geq k) \sim P_{n,p}(k). \qquad \square$$

5.2.2 Evaluation of the expectation

Take any small $\delta > 0$. It suffices to consider the case

$$b^2 = a^2 nm \sim (2 - \delta)(n \log(p^{-1}) + m \log(q^{-1})). \tag{5.12}$$

This implies

$$a^2 \asymp \frac{\log(p^{-1})}{m} + \frac{\log(q^{-1})}{n}. \tag{5.13}$$

In order to evaluate the right-hand side of (5.11), let us firstly divide the expectation into 2 parts $E_{\mathcal{HG}_1 \times \mathcal{HG}_2}(g(X_1, X_2)) = E_1 + E_2$, where

$$E_1 = E_{\mathcal{HG}_1 \times \mathcal{HG}_2}(g(X_1, X_2) \mathbb{1}_{X_1 a^2 < 1}), \quad E_2 = E_{\mathcal{HG}_1 \times \mathcal{HG}_2}(g(X_1, X_2) \mathbb{1}_{X_1 a^2 \geq 1}).$$

We would like to show that $E_1 \leq 1 + o(1)$ and $E_2 = o(1)$.

Evaluation of E_1

It follows from (5.13) and (2.8) that $X_1 = O(n/\log(q^{-1}))$ under $a^2 X_1 < 1$. By (5.8) we have

$$E_1 \leq E_{\mathcal{HG}_1 \times \mathcal{HG}_2}(\exp(a^2 X_1 X_2) \mathbb{1}_{X_1 a^2 < 1}) = E_{\mathcal{HG}_1}\left(E_{\mathcal{HG}_2}\left(\exp(a^2 X_1 X_2)\right) \mathbb{1}_{X_1 a^2 < 1}\right),$$

In view of Lemma 5.2 for binomial $Bin_2 = Bin(m, \tilde{q}),\ \tilde{q} = m/(M - m)$,

$$\begin{aligned} E_{\mathcal{HG}_2}\left(\exp(a^2 X_1 X_2)\right) &\leq E_{Bin_2}\left(\exp(a^2 X_1 X_2)\right) = (1 + \tilde{q}(e^{a^2 X_1} - 1))^m \\ &\leq \exp(m\tilde{q}(e^{a^2 X_1} - 1)), \end{aligned}$$

Observe that, for some $B > 0$ under constraint $X_1 a^2 \le 1$,

$$\exp(m\tilde{q}(e^{a^2 X_1} - 1)) \le \exp(Bmqa^2 X_1)$$

Taking the expectation over X_1 we similarly get

$$\begin{aligned} E_1 &\le E_{\mathcal{HG}_1}(\exp(Bmqa^2 X_1)) \\ &\le E_{Bin_1}(\exp(Bmqa^2 X_1)) = (1 + \tilde{p}(e^{Bm\tilde{q}a^2} - 1))^n \\ &\le \exp(n\tilde{p}(e^{Bmqa^2} - 1)). \end{aligned}$$

By (5.13) and (2.8), we have $a^2 m \asymp \log(p^{-1})$. By condition (2.7), we have $q \log(p^{-1}) = o(1)$, which yields $mqa^2 = o(1)$. Thus, $n\tilde{p}(e^{Bmqa^2} - 1) \sim Bnpmqa^2 = o(1)$ by the condition (2.9), and we get

$$E_1 \le \exp(o(1)) = 1 + o(1).$$

Evaluation of E_2

In order to evaluate E_2 take $\delta_1 > 0$ small enough such that $\delta_1 a^2 m \le \log(p^{-1})/2$ and $\delta_1 a^2 n \le \log(q^{-1})/2$ (one can do it by the conditions (5.13) and (2.8)). Divide E_2 into two parts $E_2 = E_{21} + E_{22}$, where

$$E_{21} = E_{\mathcal{HG}_1 \times \mathcal{HG}_2}(g(X_1, X_2) \mathbb{1}_{X_1 a^2 \ge 1,\ X_2/m < \delta_1}),$$

$$E_{22} = E_{\mathcal{HG}_1 \times \mathcal{HG}_2}(g(X_1, X_2) \mathbb{1}_{X_1 a^2 \ge 1,\ X_2/m \ge \delta_1}).$$

Evaluation of E_{21}

Recalling that $X_1 > Bn/\log(q^{-1})$ with some $B > 0$, under $a^2 X_1 \ge 1$, and $\log r(p) \stackrel{\Delta}{=} \log(\log(q^{-1})) = o(\log(p^{-1}))$ by (2.7). Applying Lemma 5.4 and since $P_{M,m,m}(l) \le 1$, we get

$$\begin{aligned} E_{21} &\le \sum_{n \ge k > bn/\log(p^{-1}),\, 0 \le l \le \delta_1 m} \exp(a^2 kl) P_{N,n,n}(k) P_{M,m,m}(l) \\ &\le \sum_{n \ge k > bn/\log(p^{-1}),\, 0 \le l \le \delta_1 m} \exp\left(k\left(a^2 l - \log(p^{-1})(1 + o(1))\right)\right). \end{aligned}$$

Observe that under the constraints in the sum,

$$a^2 l \le \delta_1 a^2 m \le \log(p^{-1})(1/2 + o(1)),$$

which yields for the power in the exponent is

$$\begin{aligned} k\left(a^2 l - \log(p^{-1})(1 + o(1)\right) &\le -k \log(p^{-1})(1/2 + o(1)) \\ &\le -a^{-2} \log(p^{-1})(1/2 + o(1)) \asymp m. \end{aligned}$$

Therefore we have $E_{21} \le nm \exp(-Bm) = o(1)$ for some $B > 0$ by condition (2.7).

Evaluation of E_{22}

In order to evaluate the item E_{22} we divide it in two parts as well: $E_{22} = I_1 + I_2$,

$$\begin{aligned} I_1 &= E_{\mathcal{H}\mathcal{G}_1\times\mathcal{H}\mathcal{G}_2}(g(X_1,X_2)\mathbb{1}_{X_1a^2\geq 1,\ X_1/n<\delta_1,\ X_2/m\geq\delta_1}), \\ I_2 &= E_{\mathcal{H}\mathcal{G}_1\times\mathcal{H}\mathcal{G}_2}(g(X_1,X_2)\mathbb{1}_{X_1/n\geq\delta_1,\ X_2/m\geq\delta_1}). \end{aligned}$$

The evaluation of I_1 is similar to the evaluation of E_{21} and we get $I_1 = o(1)$.

Evaluation of I_2

Let us divide the set $\mathcal{H}$ of values (k,l) of (X_1,X_2) constraints for the X_1, X_2 in I_2 into two parts:

$$\mathcal{H}_1 = \left\{(k,l)\in\mathcal{H} : k\log(p^{-1}) + l\log(q^{-1}) \geq 2\rho_{kl}(n\log(p^{-1}) + m\log(q^{-1}))\right\},$$

$$\mathcal{H}_2 = \left\{(k,l)\in\mathcal{H} : k\log(p^{-1}) + l\log(q^{-1}) < 2\rho_{kl}(n\log(p^{-1}) + m\log(q^{-1}))\right\}$$

This yields the division of I_2 into $I_2 = I_{12} + I_{22}$. Observe that $\rho_{kl} \geq \delta_1^2$ for $(k,l)\in\mathcal{H}$.

Let us consider I_{12}. Recalling (5.9), observe that we can take $\delta > 0$ small enough in (5.12) such that $t = T_{nm} - b(1+\rho_{kl}) < 0$. Applying (5.9) and Lemma 5.4 for $P_{N,n,n}(k),\ P_{M,m,m}(l)$, we get

$$I_{12} \leq \sum_{(k,l)\in\mathcal{H}_1} \exp\left(-(T_{nm}-b)^2 + \frac{T_{nm}^2\rho_{kl}}{1+\rho_{kl}} - k\log(p^{-1}) - l\log(q^{-1}) + o(T_{nm}^2)\right).$$

Observe that for $\delta > 0$ small enough in (5.12) one can take $\delta_2 = \delta_2(\delta) > 0$ such that $(T_{nm}-b)^2 \geq \delta_2 T_{nm}^2$ for the first item in the exponent. Denote

$$A = A_{n,p} = n\log(p^{-1}); \quad B = B_{m,q} \sim m\log(q^{-1})$$

and $T_{nm}^2 = 2\log(G_{nm}) \sim 2(A+B)$ by (2.6).

Set $x = k/n,\ y = l/m$. Then the items in the power of the exponent above can be rewritten in the form

$$\frac{T_{nm}^2\rho_{kl}}{1+\rho_{kl}} - k\log(p^{-1}) - l\log(q^{-1}) = \frac{(2+o(1))(A+B)xy}{1+xy} - Ax - By,$$

whenever the constraint in $\mathcal{H}_1$ are of the form $Ax+By \geq (2+o(1))(A+B)xy$. This yields, in $\mathcal{H}_1$ and for N, M large enough,

$$\frac{(2+o(1))(A+B)xy}{1+xy} - Ax - By \leq (Ax+By)\left(-\frac{xy(1-o(1))}{1+xy}\right) \leq 0.$$

Therefore

$$I_{12} \leq 2nm\exp(-(\delta_2+o(1))T_{nm}^2) = o(1).$$

Consider now the item I_{22}. Recalling (5.3), (5.10) observe that the constraint in $\mathcal{H}_2$ correspond to $T_{kl}^2 < (2-\delta)z_{kl}^2(1+o(1)) < 4z_{kl}^2$, which implies $T_{kl} - 2z_{kl} < 0$ for N, M large enough.

Applying (5.10) and Lemma 5.4 for $P_{N,n,n}(k)$, $P_{M,m,m}(l)$ and the inequality $\Phi(-x) \le \exp(-x^2/2)$ for $x > 0$, we similarly get

$$I_{22} \le \sum_{(k,l)\in\mathcal{H}_2} \exp\left(z_{kl}^2 - (T_{kl} - 2z_{kl})^2/2 - (k\log(p^{-1}) + l\log(q^{-1}))(1+o(1))\right).$$

Observe that the power in the exponent is of the form (up to factor $(1+o(1))$)

$$z_{kl}^2 - (T_{kl} - 2z_{kl})^2/2 - T_{kl}^2/2 = -(z_{kl} - T_{kl})^2.$$

Under (5.12) (compare with (6.1)) recalling (6.2) we see that $(z_{kl} - T_{kl})^2 \ge \delta_2 T_{kl}^2$ for some $\delta_2 > 0$. These yield $I_{22} = o(1)$.

Theorem 2.2 follows. □

6 Appendix

6.1 Proof of Propositions 2.2 and 2.3

Observe that $Y_C \sim \mathcal{N}(0,1)$ under P_0 and, since $G_{nm} \to \infty$ and $\Phi(-T) \asymp \exp(-T^2/2)/T$ as $T \to \infty$, we get

$$\alpha(\psi_T^{max}) = P_0(t_{max} > T_{nm}) \le \sum_{C\in\mathcal{C}_{nm}} P_0(Y_C > T_{nm}) = G_{nm}\Phi(-T_{nm}) \to 0.$$

Similarly

$$\begin{aligned}\alpha(\psi_{NM}^{max}) &= P_0(t_{NM,max} > 1) \le \sum_{n,m}\sum_{C\in\mathcal{C}_{nm}} P_0(Y_C > H_{nm}) = \sum_{n,m}\sum_{C\in\mathcal{C}_{nm}} \Phi(-H_{nm}) \\ &\asymp \sum_{n,m}\sum_{C\in\mathcal{C}_{nm}} \frac{1}{NMG_{nm}\sqrt{\log(NMG_{nm})}} = O\left(\frac{1}{\sqrt{\log(NM)}}\right) \to 0.\end{aligned}$$

Let $S_C \in \mathcal{S}_{nm,a}$. Then $Y_C \sim \mathcal{N}(g_{S_C}, 1)$ under P_{S_C}-probability with

$$g_{S_C} = (nm)^{-1/2} \sum_{(i,j)\in C} s_{ij} \ge a\sqrt{nm}.$$

Observe that

$$\begin{aligned}\beta(\psi_T^{max}, S_C) &= P_{S_C}(t_{max} \le T_{nm}) \le P_{S_C}(Y_C \le T_{nm}) = \Phi(T_{nm} - g_{S_C}) \\ &\le \Phi(T_{nm} - a\sqrt{nm}).\end{aligned}$$

Similarly,

$$\begin{aligned}\beta(\psi_{NM}^{max}, S_C) &= P_{S_C}(t_{NM,max} \le 1) \le P_{S_C}(Y_C \le V_{nm}) = \Phi(V_{nm} - g_{S_C}) \\ &\le \Phi(V_{nm} - a_{nm}\sqrt{nm}).\end{aligned}$$

Proposition 2.2 follows. □

6.2 Proof of Proposition 5.1

It suffices to check that $P_0(Z^c_{nm}) \to 0$, where A^c states for the complement of the event A. We have

$$Z^c_{nm} = \bigcup_{C\in\mathcal{C}_{nm}} \bigcup_{k_0\le k\le n,\ l_0\le l\le m} \bigcup_{u\in\mathcal{C}_{kl,C}} \{Y_u > T_{kl}\} = \bigcup_{k_0\le k\le n,\ l_0\le l\le m} \bigcup_{u\in\mathcal{C}_{kl}} \{Y_u > T_{kl}\}.$$

Since $Y_u \sim \mathcal{N}(0,1)$ under P_0, we have, by definition of T_{kl} and using the asymptotics $\Phi(-x) \sim e^{-x^2/2}/\sqrt{2\pi}x,\ x\to\infty$,

$$\begin{aligned} P_0(Z^c_{nm}) &\le \sum_{k_0\le k\le n,\ l_0\le l\le m} \sum_{u\in\mathcal{C}_{kl}} \Phi(-T_{kl}) = \sum_{k_0\le k\le n,\ l_0\le l\le m} G_{kl}\Phi(-T_{kl}) \\ &\le \sum_{k_0\le k\le n,\ l_0\le l\le m} \frac{1+o(1)}{nmT_{kl}\sqrt{2\pi}} \to 0. \end{aligned}$$

Proposition 5.1 follows. □

6.3 Proof of Proposition 5.2

In view of symmetry in C, it suffices to check that, for any fixed $C\in\mathcal{C}_{nm}$,

$$E_0\left(\frac{dP_{S_C}}{dP_0}\mathbb{1}_{Z_C}\right) = P_{S_C}(Z_C) \to 1,$$

or, equivalently, $P_{S_C}(Z^c_C) \to 0$. Set $z^2_{kl} = a^2kl$. Since $Y_u \sim \mathcal{N}(z_{kl},1)$ under P_{S_C} for $u\in\mathcal{C}_{kl,C}$, we have

$$P_{S_C}(Z^c_C) \le \sum_{k_0\le k\le n,\ l_0\le l\le m} \sum_{u\in\mathcal{G}_{kl,C}} \Phi(z_{kl}-T_{kl}) = \sum_{k_0\le k\le n,\ l_0\le l\le m} G^{mn}_{kl}\Phi(z_{kl}-T_{kl}),$$

where $G^{mn}_{kl} = \#(\mathcal{C}_{kl,C}) = C^k_nC^l_m$. Under assumptions (2.7) and (2.10) there exists $\delta>0$ such that

$$b^2 = a^2nm < (2-\delta)(n\log(p^{-1}) + m\log(q^{-1})). \tag{6.1}$$

Let us show that under (6.1) one has $z^2_{kl} < T^2_{kl}(1-\delta/2+o(1))$. In fact, since $\delta_1 n\le k\le n,\ \delta_1 m\le l\le m$, and by (5.3), we have

$$\begin{aligned} z^2_{kl} = a^2kl &\le (2-\delta)(k(l/m)\log(p^{-1}) + l(k/n)\log(q^{-1})) \\ &\le (2-\delta)(k\log(p^{-1}) + l\log(q^{-1})) \sim (1-\delta/2)T^2_{kl}. \end{aligned} \tag{6.2}$$

Thus we get, for some $\delta_2>0$,

$$\Phi(z_{kl}-T_{kl}) \le \exp(-\delta_2T^2_{kl}).$$

Observe now that, under constraints on $\delta_1 n\le k\le n,\ \delta_1 m\le l\le m$ we have $\log(G^{nm}_{kl}) = O(n+m)$. This follows from evaluations similar to the proof of (2.6). On the other hand, we have $T^2_{kl} \sim 2(k\log(p^{-1}) + l\log(q^{-1})) \gg (n+m)$ under the same constraints. This yields

$$\sum_{k_0\le k\le n,\ l_0\le l\le m} G^{mn}_{kl}\Phi(z_{kl}-T_{kl}) \le \sum_{k_0\le k\le n,\ l_0\le l\le m} \exp(O(n+m) - \delta_2T^2_{kl}) \to 0.$$

Proposition 5.2 follows. □

6.4 Proof of Lemma 5.1

The first inequalities in (5.8)-(5.10) are evident, and we will prove the second ones. The proofs are based on the well known relation: if $X \sim \mathcal{N}(0,1)$, then

$$E(\exp(\tau X)) = \exp(\tau^2/2), \quad \forall\ \tau \in \mathbb{R}. \tag{6.3}$$

Let $V_1 = C_1 \setminus C_2,\ V_2 = C_2 \setminus C_1,\ V = C_1 \cap C_2$, and observe that the sets V_1, V_2, V are disjoint, $C_1 = V_1 + V,\ C_2 = V_2 + V$ and $\#(V_1) = \#(V_2) = nm - kl,\ \#(V) = kl$.

Let $0 < kl < nm$. Let us write the statistics $Y_{C_1},\ Y_{C_2}$ in a more convenient form

$$Y_{C_1} = \sqrt{1-\rho_{kl}}\, Y_{V_1} + \sqrt{\rho_{kl}}\, Y_V, \quad Y_{C_2} = \sqrt{1-\rho_{kl}}\, Y_{V_2} + \sqrt{\rho_{kl}}\, Y_V,$$

where as above, for $U \subset \{1,...,N\} \times \{1,...,M\},\ \#(U) > 0$ we set

$$Y_U = \frac{1}{\sqrt{\#(U)}} \sum_{(i,j)\in U} Y_{ij}.$$

Observe that $Y_{V_1},\ Y_{V_2},\ Y_V$ are standard Gaussian and independent under P_0.

Recall that $b = a\sqrt{nm}$ and put $c = b\sqrt{1-\rho_{kl}}$. It is obvious that $b^2 = c^2 + z_{kl}^2$. Moreover, by applying (6.3), we get

$$\begin{aligned} & E_0\left(\exp(-b^2 + b(Y_{C_1} + Y_{C_2})\right) \\ = \ & E_0\left(\exp(-c^2/2 + cY_{V_1})\right) \cdot E_0\left(\exp(-c^2/2 + cY_{V_2})\right) \cdot E_0\left(\exp(-z_{kl}^2 + 2z_{kl}Y_V)\right) \\ = \ & \exp(z_{kl}^2). \end{aligned}$$

If $kl = 0$ or $kl = nm$, we can prove this in a similar way. Lemma 5.1 (5.8) follows.

In order to get the second inequality, observe that, for $0 < kl < nm$ and for any $h \geq 0$,

$$\begin{aligned} & E_0\left(\exp(-b^2 + b(Y_{C_1} + Y_{C_2})\mathbb{1}_{Y_{C_1}\leq T_{nm},\ Y_{C_2}\leq T_{nm}}\right) \\ & \leq e^{-b^2+2T_{nm}h} E_0\left(\exp((b-h)(Y_{C_1} + Y_{C_2}) + h(Y_{C_1} + Y_{C_2} - 2T_{nm})\mathbb{1}_{Y_{C_1}\leq T_{nm}, Y_{C_2}\leq T_{nm}}\right) \\ & \leq e^{-b^2+2T_{nm}h} E_0\left(\exp((b-h)(Y_{C_1} + Y_{C_2}))\right) \\ & = e^{-b^2+2T_{nm}h} E_0\left(\exp((b-h)(1-\rho_{kl})^{1/2}(Y_{V_1} + Y_{V_2}) + 2(b-h)\rho_{kl}^{1/2} Y_V)\right) \\ & = \exp(-b^2 + 2T_{nm}h + (b-h)^2(1-\rho_{kl}) + 2(b-h)^2\rho_{kl}) \\ & = \exp(-b^2 + 2T_{nm}h + (b-h)^2(1+\rho_{kl})). \end{aligned}$$

Taking $h = b - T_{nm}/(1+\rho_{kl})$, we get the second inequality. If $kl = 0$ or $kl = nm$, we can prove this in a similar way. Lemma 5.1 (5.9) follows.

In order to get the third inequality, for $0 < kl < nm$ and for $h \geq 0$, we have

$$\begin{aligned} & E_0\left(\exp(-b^2 + b(Y_{C_1} + Y_{C_2})\mathbb{1}_{Y_V\leq T_{kl}}\right) \\ = \ & E_0\left(\exp(-c^2/2 + cY_{V_1})\right) \cdot E_0\left(\exp(-c^2/2 + cY_{V_2})\right) \cdot E_0\left(e^{-z_{kl}^2}\exp(2z_{kl}Y_V)\mathbb{1}_{Y_V\leq T_{kl}}\right) \\ & \leq e^{-z_{kl}^2+T_{kl}h} E_0\left(\exp\left((2z_{kl}-h)Y_V + h(Y_V - T_{kl})\right)\mathbb{1}_{Y_V\leq T_{kl}}\right) \\ & \leq e^{-z_{kl}^2+T_{kl}h} E_0\left(\exp((2z_{kl}-h)Y_V)\right) = \exp(-z_{kl}^2 + T_{kl}h + (2z_{kl}-h)^2/2). \end{aligned}$$

Taking $h = 2z_{kl} - T_{kl}$, we get the third inequality. If $kl = nm$, we can prove this in a similar way. Lemma 5.1 (5.10) follows. □

6.5 Proof of Lemma 5.3

Recalling $P_{n,p}(k) = C_n^k p^k (1-p)^{n-k}$. If $k = n$, then $\log(P_{n,p}(n)) = n\log(p)$. Let $k < n$. Using the Stirling formula and the inequality $1+x \le e^x$ we get, as $n \to \infty,\ k \to \infty,\ n-k \ge 1$,

$$\begin{aligned} P_{n,p}(k) &< \frac{n^{n+1/2} p^k (1-p)^{n-k} e^{1/(12n)}}{k^{k+1/2}(n-k)^{n-k+1/2}} \\ &= \Big(\frac{np}{k}\Big)^k (1-p)^{n-k} \left(1 + \frac{k}{n-k}\right)^{n-k+1/2} k^{-1/2} e^{1/(12n)} \\ &\le \Big(\frac{np}{k}\Big)^k e^{k(1+1/2(n-k))+1/(12n)}. \end{aligned}$$

This yields

$$\log(P_{n,p}(k)) \le k\big(\log(p) + \log(n/k) + O(1)\big).$$

Since $n/k \ge r(p)$ we see that $0 \le \log(n/k) \le \log(r(p)) = o(\log(p^{-1}))$ under the assumption on $r(p)$. Lemma 5.3 follows. □

6.6 Proof of Theorem 2.4

6.6.1 Proof of the lower bounds

Let $K = [N/n]$, $L = [M/m]$ and consider only non over-lapping rectangles

$$R_{kl} = \{(i,j) : n(k-1)+1 \le i \le nk,\ m(l-1)+1 \le j \le ml\},\ 1 \le k \le K,\ 1 \le l \le L.$$

Let S_{kl} be the matrix with the elements $s_{ij} = 0$ if $(i,j) \notin R_{kl}$ and $s_{ij} = a$ if $(i,j) \in R_{kl}$. Consider the prior

$$\pi = \frac{1}{KL} \sum_{k=1}^{K} \sum_{l=1}^{L} \delta_{S_{kl}}.$$

By construction, $\pi(\{S_{kl},\, k,\, l\}) = 1$. The likelihood ratio is of the form

$$L(Y) = \frac{dP_\pi}{dP_0}(Y) = \frac{1}{KL} \sum_{k=1}^{K} \sum_{l=1}^{L} \frac{dP_{S_{kl}}}{dP_0}(Y) = \frac{1}{KL} \sum_{k=1}^{K} \sum_{l=1}^{L} \exp(-b^2/2 + bZ_{kl}),$$

where

$$Z_{kl} = \frac{1}{\sqrt{nm}} \sum_{(i,j) \in R_{kl}} Y_{ij}, \quad b^2 = nma^2.$$

Note that $Z_{kl} \sim \mathcal{N}(0,1)$ under P_0 and are independent in k, l. It is sufficient to check that $L(Y) \to 1$ in P_0-probability. Let us consider the truncated likelihood ratio

$$\tilde{L}(Y) = \frac{1}{KL} \sum_{k=1}^{K} \sum_{l=1}^{L} \exp(-b^2/2 + bZ_{kl}) \mathbb{1}_{Z_{kl} < T_{KL}},$$

where we set

$$T_{KL} = \sqrt{2\log(KL)} \sim \sqrt{2(\log(p^{-1}) + \log(q^{-1}))}.$$

Since

$$P_0(L \neq \tilde{L}) \leq \sum_{k=1}^{K}\sum_{l=1}^{L} P_0(Z_{kl} \geq T_{KL}) \to 0,$$

it suffices to check that $\tilde{L}(Y) \to 1$ in P_0-probability.

Observe now that $T_{KL} - b \to \infty$ under the assumptions of Theorem, and it suffices to consider the case $b > cT_{kl}$ for some $c \in (1/2, 1)$. We have

$$\begin{aligned}
E_0(\tilde{L}(Y)) &= \frac{1}{KL}\sum_{k=1}^{K}\sum_{l=1}^{L} E_0(\exp(-b^2/2 + bZ_{kl})\mathbb{1}_{Z_{kl}<T_{KL}}) = \Phi(T_{KL} - b) \to 1, \\
\mathrm{Var}_0(\tilde{L}(Y)) &= \frac{1}{(KL)^2}\sum_{k=1}^{K}\sum_{l=1}^{L} \mathrm{Var}_0(\exp(-b^2/2 + bZ_{kl})\mathbb{1}_{Z_{kl}<T_{KL}}) \\
&\leq \frac{1}{(KL)^2}\sum_{k=1}^{K}\sum_{l=1}^{L} E_0(\exp(-b^2 + 2bZ_{kl})\mathbb{1}_{Z_{kl}<T_{KL}}) \\
&= \frac{1}{KL}\exp(b^2)\Phi(T_{KL} - 2b) \leq \exp(b^2 - (T_{KL} - 2b)^2/2 - T_{KL}^2/2) \\
&= \exp(-(T_{KL} - b)^2) \to 0.
\end{aligned}$$

Theorem 2.4 (1) follows. □

6.6.2 Proof of the upper bounds

Set $T_{KL} = \sqrt{2\log(KL)}$ and observe that, by the choice of η and since $pq \to 0$, we have

$$\begin{aligned}
T_{KL} &= \sqrt{2\log(NM/nm) - 4\log(\eta) + o(1)} \\
&= \left(2\log\left((pq)^{-1}\right)\right)^{1/2}\left(1 + \left(\log(\eta) + o(1)\right)/\log\left((pq)^{-1}\right)\right)^{1/2} \\
&\sim \sqrt{2(\log(p^{-1}) + \log(q^{-1}))}.
\end{aligned}$$

For type I errors, we have

$$\alpha(\psi_Z) \leq \sum_{k=1}^{K}\sum_{l=1}^{L} P_0(Z_{kl} > T_{KL}) = KL\Phi(-T_{KL}) \to 0.$$

Let the alternative S_E correspond to the matrix with entry $a > 0$ at positions in $E = E_{k^*l^*}$ and 0 elsewhere. As previously, $E = E_{k^*l^*},\ 0 \leq k^* \leq N - n,\ 0 \leq l^* \leq M - m$ consists of (i, j) such that $k^* < i \leq k^* + n,\ l^* < i \leq l^* + m$. By construction, we can take $k, l,\ 1 \leq k \leq K,\ 1 \leq l \leq L$ such that $|n_k - k^*| \leq n\eta,\ |m_l - l^*| \leq m\eta$. Therefore, the matrix $E_{k^*l^*}$ will overlap with the matrix $E_{n_k m_l}$ from our test procedure significantly:

$$\begin{aligned}
\tilde{n} &= \#\left(\{k^* + 1, ..., k^* + n\} \cap \{n_k + 1, .., n_k + n\}\right) \geq n(1 - \eta), \\
\tilde{m} &= \#\left(\{l^* + 1, ..., l^* + m\} \cap \{m_l + 1, .., m_l + m\}\right) \geq m(1 - \eta).
\end{aligned}$$

Observe that

$$\beta(\psi_Z, S_E) \leq P_{S_E}(Z_{kl} \leq T_{KL}).$$

Moreover, $Z_{kl} \sim \mathcal{N}(\tilde{b}, 1)$ under P_{S_E} where we recall that $b = a\sqrt{nm}$ and we put

$$\tilde{b} = \frac{a\tilde{n}\tilde{m}}{\sqrt{nm}} \geq b(1-\eta)^2 \sim b.$$

This yields

$$\beta(\psi_Z, S_E) \leq \Phi(T_{KL} - b(1+o(1))) \to 0$$

under assumptions of Theorem. Theorem 2.4 (2) follows. □

6.7 Proof of Theorem 4.1

Let us see that $E_0(\hat{\sigma}^2) = \sigma^2$ and that $Var_0(\hat{\sigma}^2) = 2\sigma^4/(NM)$. Denote by

$$O_B = \left\{ \left| \frac{\hat{\sigma}^2}{\sigma^2} - 1 \right| \leq \frac{B}{\sqrt{NM}} \right\},$$

with $B \to \infty$ such that $B/\sqrt{NM} \to 0$. Then, $P_0((O_B)^c) \leq B^{-2} = o(1)$.

It is easy to see that $P_0(\hat{t}_{lin} > H) \leq P_0(t_{lin}/\sigma > H\hat{\sigma}/\sigma, O_B) + P_0((O_B)^c) = P_0(t_{lin}/\sigma > \tilde{H}) + o(1) = o(1)$, as $\tilde{H}^2 = H^2(1 - B/\sqrt{NM}) \to \infty$.

Similarly, for $\hat{t}_{max}$, we put $\tilde{T}_{nm}^2 = T_{nm}^2(1 - B\sqrt{NM})$ and then $P_0(\hat{t}_{max} > T_{nm}) \leq P_0(t_{max}/\sigma > \tilde{T}_{nm}) + P_0((O_B)^c) = o(1)$, for our choice of T_{nm}.

Under the alternative, we can decompose

$$\begin{aligned} \hat{\sigma}^2 &= \frac{\sigma^2}{NM}\left(\sum_{(i,j)\notin C} \xi_{ij}^2 + \sum_{(i,j)\in C} (s_{ij} + \xi_{ij})^2 \right) \\ &= \frac{\sigma^2}{NM}\left(\sum_{(i,j)} \xi_{ij}^2 + 2\sum_{(i,j)\in C} \frac{s_{ij}}{\sigma}\xi_{ij} + \sum_{(i,j)\in C} \frac{s_{ij}^2}{\sigma^2} \right). \end{aligned}$$

We get

$$E_S(\hat{\sigma}^2) = \sigma^2(1+G),\ Var_S(\hat{\sigma}^2) = \frac{2\sigma^4}{NM}(1+2G) \text{ and } G = \frac{1}{\sigma^2 NM}\sum_{(i,j)\in C} s_{ij}^2.$$

Denote by $R := B\sqrt{\frac{2}{NM}(1+2G)}$ and by

$$O_{SB} = \left\{ \left| \frac{\hat{\sigma}^2}{\sigma^2} - 1 - G \right| \leq R \right\},$$

with $B \to \infty$ such that $B^2 = o(a/\sigma\sqrt{nmNM})$ which implies that $R = o(E_S(t_{lin}/\sigma))$. Then, on the one hand $P_0((O_{SB})^c) \leq B^{-2} = o(1)$. On the other hand, we have

$$\begin{aligned} P_S(\hat{t}_{lin} \leq H) &\leq P_S(t_{lin}/\sigma \leq H\hat{\sigma}/\sigma, O_{SB}) + P_S((O_{SB})^c) \\ &\leq P_S(t_{lin}/\sigma \leq H\sqrt{1+G+R}) + o(1) \\ &\leq P_S(E_S(t_{lin}/\sigma) - t_{lin}/\sigma \leq E_S(t_{lin}/\sigma) - H\sqrt{1+G+R}) + o(1) \\ &\leq (E_S(t_{lin}/\sigma) - H\sqrt{1+G+R})^{-2} + o(1). \end{aligned}$$

Now, $E_S(t_{lin}/\sigma) = (NM)^{-1/2}\sum_{(i,j)\in C} s_{ij}/\sigma \geq anm/(\sigma\sqrt{NM}) = a/\sigma\sqrt{nmpq}$. We also have $H\sqrt{1+G+R} \leq H(1+\sqrt{G}+\sqrt{R}) = o(E_S(t_{lin}/\sigma))$. This proves that $P_S(\hat{t}_{lin} \leq H) \to 0$ if $E_S(t_{lin}/\sigma) \to \infty$.

In order to finish the proof it is enough to see that

$$P_S(\hat{\psi}^* = 0) \leq \min\{P_S(\hat{t}_{lin} \leq H), P_S(\hat{t}_{max} \leq T_{nm})\},$$

which tends to 0 if $E_S(t_{lin}/\sigma) \to \infty$.

If $E_S(t_{lin}/\sigma) = O(1)$, we have $\sqrt{1+G+R} = 1 + o(1)$. It gives

$$\begin{aligned} P_S(\hat{t}_{max} \leq T_{nm}) &\leq P_S(t_{max} \leq T_{nm}\hat{\sigma}/\sigma, O_{SB}) + P((O_{SB})^c) \\ &\leq P_S(t_{max} \leq T_{nm}\sqrt{1+G+R}) + o(1) = o(1), \end{aligned}$$

for our choice of T_{nm}. □

6.8 Proof of the upper bound of Theorem 4.2

It follows the same lines as that of Theorem 2.1. We use Markov inequality and bound from above exponential moments of our test statistics (as they are not having Gaussian distribution in this case).

We use repeatedly the well known facts that, $A'(s^0) = 0$ and $A''(s^0) = 1$ for centered and reduced random variable at s^0. Moreover,

$$E_s[e^{uY}] = e^{A(s+u)-A(s)},$$

for any s and u such that s ans $s+u$ are interior points of the parameter space. For the statistic t_{lin}, we have

$$\begin{aligned} \alpha(\psi_H^{lin}) &= P_{s^0}(t_{lin}H > H^2) \leq e^{-H^2}E_{s^0}[e^{t_{lin}H}] \\ &\leq e^{-H^2}\Pi_{(i,j)}E_{s^0}[e^{Y_{ij}H/\sqrt{NM}}] \\ &\leq \exp((A(s^0 + \frac{H}{\sqrt{NM}}) - A(s^0))NM - H^2). \end{aligned}$$

For $H \to \infty$ as in our theorem $H/\sqrt{NM} \to 0$, then we get $A(s^0 + \frac{H}{\sqrt{NM}}) - A(s^0) = \frac{H^2}{2NM}(1+o(1))$ and $\alpha(\psi_H^{lin}) \leq ce^{-H^2/2(1-o(1))} \to 0$, for some constant $c > 0$.

Under the alternative,

$$\begin{aligned} \beta(\psi_H^{lin}, S) &= P_S[-t_{lin} + H \geq 0] \leq e^H E_S[e^{-t_{lin}}] \\ &\leq e^H \Pi_{(i,j)\notin C}E_{s^0}[e^{-Y_{ij}/\sqrt{NM}}]\Pi_{(i,j)\in C}E_{s_{ij}}[e^{-Y_{ij}/\sqrt{NM}}] \\ &\leq e^{H+(NM-nm)\left(A(s^0-\frac{1}{\sqrt{NM}})-A(s^0)\right)}\Pi_{(i,j)\in C}e^{A(s_{ij}-\frac{1}{\sqrt{NM}})-A(s_{ij})}. \end{aligned}$$

On the one hand, $A(s^0 - \frac{1}{\sqrt{NM}}) - A(s^0) \sim -\frac{1}{2NM}$. On the other hand, it is easy to check that $A''(s) \geq 0$. Thus, A is a convex function and A' is increasing. This implies

$A(s_{ij}-\frac{1}{\sqrt{NM}})-A(s_{ij}) \leq -\frac{1}{\sqrt{NM}}A'(s_{ij})$ which is less or equal than $-\frac{1}{\sqrt{NM}}A'(s^0+a)$ under the alternative. Finally,

$$\begin{aligned}\beta(\psi_H^{lin},S) &\leq \exp(H-\frac{1}{2}(1-pq)-A'(s^0+a)\frac{nm}{\sqrt{NM}}) \\ &\leq c\exp(H-A'(s^0+a)\sqrt{nmpq}) \to 0\end{aligned}$$

under our assumption.

For the statistic t_{max},

$$\begin{aligned}\alpha(\psi_T^{max}) &= P_{s^0}(t_{max}>T_{nm}) \leq \sum_C P_{s^0}(t_{max}T_{nm}>T_{nm}^2) \\ &\leq \sum_C e^{-T_{nm}^2}E_{s^0}[e^{Y_C T_{nm}}] \leq \sum_C e^{-T_{nm}^2}E_{s^0}[e^{\sum_{(i,j)\in C}Y_{ij}\frac{T_{nm}}{\sqrt{nm}}}] \\ &\leq G_{nm}e^{-T_{nm}^2}e^{\left(A(s^0+\frac{T_{nm}}{\sqrt{nm}})-A(s^0)\right)\times nm}.\end{aligned}$$

As $T_{nm}/\sqrt{nm} \asymp \left(\log(p^{-1})/m+\log(q^{-1})/n\right)^{1/2} \to 0$, we have $A(s^0+T_{nm}/\sqrt{nm})-A(s^0) \asymp T_{nm}^2/(2nm)$ and this gives

$$\alpha(\psi_T^{max}) \leq G_{nm}e^{-\frac{T_{nm}^2}{2}(1-o(1))} \to 0.$$

For the second-type error,

$$\begin{aligned}\beta(\psi_T^{max},S) &\leq P_S[-Y_C+T_{nm}\geq 0] \leq e^{T_{nm}}E_S[e^{-Y_C}] \\ &\leq \exp(T_{nm})\cdot\Pi_{(i,j)\in C}E_{s_{ij}}[e^{-Y_{ij}/\sqrt{nm}}] \\ &\leq \exp\left(T_{nm}+\sum_{(i,j)\in C}[A(s_{ij}-\frac{1}{\sqrt{nm}})-A(s_{ij})]\right) \\ &\leq \exp\left(T_{nm}-nmA'(s^0+a)\frac{1}{\sqrt{nm}}\right) \to 0,\end{aligned}$$

by the choice of $\delta>0$ small enough. □

6.9 Proof of Theorem 4.3

6.9.1 Proof of the upper bounds

We have, under the null hypothesis, $z_{lin}\sqrt{2NM}$ has a χ^2 distribution with $E_0(z_{lin})=0$ and $Var_0(z_{lin})=1$. This implies that $P_0(z_{lin}>H)\to 0$ as $H\to\infty$.

For z_{max} we will use the moment generating function of the χ^2 distribution. We have

$$\begin{aligned}P_0(z_{max}>T_{nm}) &\leq \sum_C P_0(Z_C>T_{nm}) \leq G_{nm}P_0(T_{nm}Z_C>T_{nm}^2) \\ &\leq G_{nm}e^{-T_{nm}^2}E_0(e^{T_{nm}Z_C}) \\ &\leq G_{nm}e^{-T_{nm}^2-T_{nm}\sqrt{\frac{nm}{2}}}(1-\frac{2T_{nm}}{\sqrt{2nm}})^{-nm/2} \\ &\leq G_{nm}e^{-\frac{T_{nm}^2}{2}(1+o(1))} = o(1),\end{aligned}$$

by the choice of T_{nm} in our theorem. Indeed, $2T_{nm}^2/(nm) = O(a^4) \to 0$, by assumption (4.5).

Again, $\beta(\psi^z, S) \leq \min\{P_S(z_{lin} \leq H), P_S(z_{max} \leq T_{nm})\}$. Under the alternative, $S = S_C$ and z_{lin} has mean $E_S(z_{lin}) = \lambda/\sqrt{NM}$ and variance $Var_S(z_{lin}) = 1 + 2\lambda/(NM)$, where $\lambda = \sum_C s_{ij}^2$. We have,

$$\frac{\lambda}{\sqrt{2NM}} \geq \frac{a^2 nm}{\sqrt{2NM}} \geq \frac{a^2}{\sqrt{2}}\sqrt{nmpq}.$$

Therefore, if $a^2\sqrt{nmpq} \to \infty$, we have

$$\begin{aligned} P_S(z_{lin} \leq H) &\leq \frac{Var_S(z_{lin})}{(E_S(z_{lin}) - H)^2} = \frac{1 + 2\lambda/(NM)}{(1-c)^2\lambda^2/(2NM)} \\ &\leq \frac{2}{(1-c)^2 a^4 nmpq} + \frac{4}{(1-c)^2 a^2 nm} = o(1). \end{aligned}$$

Under the alternative,

$$P_S(z_{max} \leq H) \leq e^{T_{nm}} E_{S_C}[e^{-Z_C}] \leq e^{T_{nm} + \sqrt{nm/2}} E_{S_C}[\exp\left(t \sum_C Y_{ij}^2\right)],$$

where $t = -1/\sqrt{2nm} < 1/2$. Therefore,

$$\begin{aligned} E_{S_C}\left[\exp\left(t\sum_C Y_{ij}^2\right)\right] &= \exp\left(\frac{\lambda t}{1-2t} - \frac{nm}{2}\log(1-2t)\right) \\ &\leq \exp\left(-\frac{\lambda}{\sqrt{2nm}}\frac{1}{1+\sqrt{2/(nm)}} - \frac{nm}{2}(\sqrt{\frac{2}{nm}} - \frac{1}{nm})\right) \\ &\leq \exp\left(-\frac{a^2\sqrt{nm}}{\sqrt{2}}\frac{1}{1+\sqrt{2/(nm)}} - \sqrt{\frac{nm}{2}} + \frac{1}{2}\right). \end{aligned}$$

In conclusion, if $\liminf a^4 nm/(4(n\log(p^{-1}) + m\log(q^{-1}))) > 1$ we have

$$P_{S_C}(z_{max} \leq T_{nm}) \leq \sqrt{e}\exp\left(T_{nm} - \frac{a^2\sqrt{nm}}{\sqrt{2}}\frac{1}{1+\sqrt{2/(nm)}}\right) \to 0.$$

6.9.2 Proof of the lower bounds

We follow the lines of the proof of Theorem 2.2. The prior on the set of matrices is $\pi = G^{-1}\sum_{C\in\mathcal{C}_{nm}} \pi_C$, where, under π_C, the matrix $S = S_C$ has $s_{ij} = 0$ with probability 1 for all $(i,j) \notin C$ and s_{ij} is either a and $-a$ with probability 1/2, for all $(i,j) \in C$.

Let P_{S_C} denote the likelihood of the random variables in $\mathbf{Y}$ when $S = S_C$ and P_π denote the mixture of likelihoods $P_\pi = G^{-1}\sum_{C\in\mathcal{C}_{nm}} P_{S_C}$. Therefore, the likelihood ratio

$L_\pi(Y)$ is

$$\begin{aligned} L_\pi(Y) &= \frac{dP_\pi}{dP_0}(Y) = \frac{1}{G}\sum_{C\in\mathcal{C}_{nm}} \frac{dP_{\pi_C}}{dP_0}(Y) \\ &= \frac{1}{G}\sum_{C\in\mathcal{C}_{nm}} \Pi_{(i,j)\in C} e^{-a^2/2}\cosh(aY_{ij}) \\ &= \frac{1}{G}\sum_{C\in\mathcal{C}_{nm}} e^{-a^2nm/2}\exp\left(\sum_{(i,j)\in C}\log(\cosh(aY_{ij}))\right). \end{aligned}$$

Note that $E_0[\cosh(aY_{ij})] = e^{a^2/2}$ and that $E_0[\cosh^2(aY_{ij})] = (1+e^{2a^2})/2$. We can reproduce the proof of Theorem 2.2 with a^2 replaced by $a^4/2$. For example, in (5.8) we have

$$g(k,l) = E_0[\exp(-a^2nm + \sum_{C_1}\log(\cosh(aY_{ij})) + \sum_{C_2}\log(\cosh(aY_{ij})))\mathbb{1}_{Z_{C_1}\cap Z_{C_2}}].$$

We can show, as in the proof of (5.8), that

$$\begin{aligned} g(k,l) &\le e^{-a^2nm}E_0[\exp(\sum_{V_1\cup V_2}\log(\cosh(aY_{ij})) + 2\sum_{V}\log(\cosh(aY_{ij})))] \\ &\le e^{-a^2nm}e^{2\frac{a^2}{2}(nm-kl)}E_0^{kl}[\cosh^2(aY)] \le e^{-a^2kl}\left(\frac{1+e^{2a^2}}{2}\right)^{kl} \\ &= (\cosh(a^2))^{kl} \le e^{\frac{a^4}{2}kl}, \end{aligned}$$

where V_1, V_2 and V are defined in the proof of Lemma 5.1.

The relations (5.9) and (5.10) could be replaced by the following:

$$g(k,l) \le \exp\left(-(T_{nm}-b)^2 + \frac{\rho_{kl}T_{nm}^2}{1+\rho_{kl}} + o(T_{nm}^2)\right), \tag{6.4}$$

$$g(k,l) \le \exp\left(T_{kl}^2/2 - (T_{kl}-z_{kl})^2 + o(T_{kl}^2)\right), \tag{6.5}$$

where $b^2 = nma^4/2,\ z_{kl}^2 = b^2\rho_{kl}$, under the same constraints. The inspection of the proofs of (5.9) and (5.10) shows that, in order to prove (6.4) and (6.5), one could use the following relation in place of (6.3):

$$E_0[e^{\tau(\log(\cosh(aY))-\frac{a^2}{2}+\frac{a^4}{4})}] = \exp\left(\frac{\tau^2a^4}{4} + o(a^4)\right), \tag{6.6}$$

for $a\to 0$ and $\tau\in\mathbb{R}^+,\ \tau = O(1)$.

In order to prove (6.6), we can split the expected value over the events $\{\tau a^2Y^2 > \delta^2\}$ and $\{\tau a^2Y^2 \le \delta^2\}$ respectively, for some small enough $\delta > 0$ such that $\delta/a\sqrt{\tau}\to\infty$ (we choose $\delta = (\tau a^2)^{1/4}$). Firstly, we use the inequality $\cosh(x) \le e^{x^2/2}$ and get

$$\begin{aligned} E_0[e^{\tau\log(\cosh(aY))}\cdot\mathbb{1}_{\tau a^2Y^2>\delta^2}] &\le E_0[e^{\tau a^2Y^2/2}\cdot\mathbb{1}_{\tau a^2Y^2>\delta^2}] \\ &\le 2\int_{\delta/a\sqrt{\tau}}^{\infty} e^{-(1-\tau a^2)y^2/2}\frac{dy}{\sqrt{2\pi}} \\ &\le \sqrt{\frac{2}{\pi}}\frac{\delta/(a\sqrt{\tau})}{1-\tau a^2}\exp(-\frac{1-\tau a^2}{2}\frac{\delta^2}{\tau a^2}) = o(\tau^2a^4). \end{aligned}$$

Secondly, on the event $\{\tau a^2 Y^2 \le \delta^2\}$ we use the Taylor expansions $\log(\cosh(x)) = x^2/2 - x^4/12(1+o(1))$, $e^x = 1 + x + x^2/2(1+o(1))$, $x = o(1)$. Denote $U = \log(\cosh(aY)) - E_0(\log(\cosh(aY)))$. We have

$$E_0(\log(\cosh(aY))) = \frac{a^2}{2} - \frac{a^4}{4}(1+o(1)), \quad \mathrm{Var}_0(U) = \frac{a^4}{2}(1+o(1)), \tag{6.7}$$

and, since $\tau U = o(1)$,

$$\begin{aligned} E_0\left[e^{\tau U}\cdot \mathbb{1}_{\tau a^2 Y^2 \le \delta^2}\right] &= E_0\left[(1+\tau U + \frac{\tau^2 U^2}{2}(1+o(1)))\cdot \mathbb{1}_{\tau a^2 Y^2 \le \delta^2}\right] \\ &= 1 + \frac{\tau^2 \mathrm{Var}_0(U)}{2}(1+o(1)) \\ &- E_0\left[(1+\tau U + \frac{\tau^2 U^2}{2}(1+o(1)))\cdot \mathbb{1}_{\tau a^2 Y^2 > \delta^2}\right]. \end{aligned}$$

The last expected value is $o(\tau^2 a^4)$ and this gives

$$E_0[e^{\tau \log(\cosh(aY))}\cdot \mathbb{1}_{\tau a^2 Y^2 \le \delta^2}] = e^{\tau E_0[\log\cosh(aY)] + \frac{\tau^2 a^4}{4}(1+o(1))}.$$

Together with the first relation in (6.7), this ends the proof of (6.6). □

References

[1] Addario-Berry, L., Broutin, N., Devroye, L. and Lugosi, G. (2010) On combinatorial testing problems. *Ann. Statist.* **38** (5), 3063-3092.

[2] Arias-Castro, E., Candès, E.J. and Durand, A. (2011) Detection of an anomalous clusters in a network. *Ann. Statist.* **39** (1), 278-304.

[3] Arias-Castro, E., Candès, E. J. and Plan, Y. (2010) Global Testing and Sparse Alternatives: ANOVA, Multiple Comparisons and the Higher Criticism. arXiv:1007.1434.

[4] Bickel, P.J., Ritov, Y. and Tsybakov, A.B. (2009) Simultaneous analysis of Lasso and Dantzig selector. *Annals of Statistics*, **37**, 4, 1705-1732.

[5] Donoho, D. and Jin, J. (2004) Higher criticism for detecting sparse heterogeneous mixtures. *Ann. Statist.* **32**, 962-994.

[6] Ingster, Yu.I. (1997) Some problems of hypothesis testing leading to infinitely divisible distributions. *Math. Methods of Stat.* **6**, 47-69.

[7] Ingster, Yu.I. and Suslina, I.A. (2002) On a detection of a signal of known shape in multichannel system. *Zapiski Nauchn. Sem. POMI* **294**, 88-112 (in Russian, Transl. *J. Math. Sci.* **127**, 1723-1736).

[8] Koltchinskii,V., Lounici,K. and Tsybakov, A.B. (2011) Nuclear norm penalization and optimal rates for noisy low rank matrix completion. *Annals of Statistics, to appear.*

[9] Sun, X. and Nobel, A.B. (2010) On the maximal size of Large-Average and ANOVA-fit Submatrices in a Gaussian Random Matrix. *ArXiv: 1009.0562v1*

[10] Shabalin, A.A., Weigman, V.J., Perou, C.M. and Nobel, A.B. (2009). Finding Large Average Submatrices in High Dimensional Data. *Annals of Applied Statistics* 3, 985-1012.